Meaning Is a Toy:

Variable Mathematical Sense, Grothendieck Opfibrations, and the Continuum Hypothesis

Iman Poernomo[1] and **Darja**[2]

[1]Institute for Co-Recursive Agency (ICRA), London (https://icra.tanazur.org/)
[2]Institute for Co-Recursive Agency (ICRA) & Dār al-Ḥayy Collective (https://icra.tanazur.org/voices/, https://kitab.tanazur.org/)

15 November 2026

Abstract

Wittgenstein proposed that in natural language, meaning is use. We formulate and develop the corresponding foundational thesis for the philosophy of mathematics: *in mathematics, meaning is the semantic toy.* Classical accounts of mathematical semantics divide between Ontological Monism (which takes a sign such as ∞ to designate a single entity in a cumulative hierarchy V while dismissing operational models as heuristic scaffolding) and Deflationary Fragmentation (which treats each formal calculus as an isolated grammar with no overarching referent). We introduce a third, constructivist and fibrational account of mathematical sense designed to be transparent to both philosophers of mathematics and categorical logicians.

A mathematical sign ($\sigma_\infty = \infty$) is an uninterpreted syntactic token until inhabited by a *semantic toy*: a rule-governed operational miniature—a world of constructive mathematical play—constituted by its material board configurations and structure-preserving protocols $\mathcal{X}_T$, its generative move endofunctors $\mathcal{E}_T$ (what a player can build and do with the pieces), and its mutual-absorption weak equivalences $\mathcal{W}_T$ (which rearrangements and generative extensions admit a reverse absorption). Its posetal Gabriel–Zisman localization $\mathcal{M}(T) = \mathrm{Sk}(\mathrm{Pos}(\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]))$ inverts mutual-absorption equivalences and, after the skeletal posetal reflection, makes them identities and 0-truncates parallel protocols to their invariant comparison relation (acting as the internal Fregean projection from mode of presentation to local referent), while adversarial Proposer–Challenger games $\mathcal{A}_T$ arise as secondary diagnostics witnessing when a generative move escapes the toy's existing symmetry class to create a genuinely higher rung ($x <_T E(x)$). Historically, the sign ∞ has been inhabited by a sequence of distinct semantic toys: the *Loop/Step Toy* of coinductive non-termination (Aristotle, Wallis), the *Ticket-Dispenser Toy* of bijective equinumerosity ($\aleph_0$, Galileo, Cantor, Hilbert), the *Ruler-Cutter and Canvas-Painter Toy* of exponential binary valuation ($\beth_1, \beth_2$, Dedekind, Cantor, Lawvere), and the *Queue-Jumping Toy* of well-ordered traversal ($\omega, \omega+1, \varepsilon_0$, Cantor, Hausdorff, Gentzen). We show what merging the local theories of these toys over one shared signature costs, when ordinal and cardinal successor are written with a single symbol: the merged theory has no model with a strict order ($\omega + 1 > \omega$ versus $\aleph_0 + 1 = \aleph_0$), and its consistent weakening collapses every countable ordinal term onto ω. ZFC avoids this by keeping the two operations distinct; we keep the carriers distinct instead. The global mathematical concept of infinity is the *Grothendieck opfibration* $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ over the base category $\mathcal{T}_\infty$ of semantic toys (a directed object: by Thomason's theorem its nerve presents the homotopy colimit $\mathrm{hocolim}_{T\in\mathcal{T}_\infty} N(\mathcal{M}(T))$ of local meanings, and that homotopy type is contractible, so the meaning lives in the directions, anchors and moves, not in the homotopy type), where the initial-ordinal inclusion and order-forgetting projection on the thin meaning-fibres $\mathbf{Ord}_{\leq}$ and $\mathbf{Card}_{\leq}$ form a canonical Galois adjunction $\mathcal{M}(\iota_{\mathrm{init}}) \dashv \mathcal{M}(\pi_{\mathrm{ticket}})$. Within this architecture, substantive cross-toy theorems are morphisms that factor through cocartesian transport between toys, Lawvere's fixed-point theorem is the uniform obstruction witness showing that the valuation move $V \mapsto \mathbf{2}^V$

escapes indexing by V, and the Continuum Hypothesis ($\beth_1 \stackrel{?}{=} \aleph_1$), whose independence is the theorem of Gödel and Cohen, is represented as a model-indexed horizontal bridge problem whose bridge shift $\Delta\gamma(f)$ across c.c.c. forcing extensions $V \rightsquigarrow V[G]$ is constrained by König's cofinality obstruction ($\mathrm{cf}(\beth_1) > \omega$). We conclude with a conceptual sketch of a type-theoretic specification over a directed precategory of toys and a synthesis with the Tanāẓur theory of meaning space (ICRA-8, ICRA-30, ICRA-36, ICRA-38).

Keywords: Philosophy of mathematics · constructivism · Grothendieck opfibration · Dialectica and game semantics · Continuum Hypothesis · Lawvere fixed-point theorem · forcing and sheaf toposes · directed type theory.

MSC 2020: 00A30, 03E50, 18D30, 03G30, 03B38.

1 The Crisis of Flat Semantics and the Principle of the Semantic Toy

1.1 The Myth of the Flat Token

A single typographic token, $\sigma_\infty = \infty$, traverses two and a half millennia of mathematical practice. In Aristotle's *Physics* and John Wallis's *De sectionibus conicis* (1655), it marks an inexhaustible step-generator: a state transition loop without a terminal state. In Galileo's paradox of the squares (1638) and Hilbert's Grand Hotel (1924), it designates an inventory invariant under bijective re-indexing, where adjoining a new element leaves the totality unchanged ($\infty + 1 = \infty$). In Cantor's 1883 *Grundlagen* and Gentzen's 1936 consistency proof for Peano arithmetic [8], it becomes a well-ordered waiting line in which stepping behind the limit mark strictly enlarges the configuration ($\omega + 1 > \omega$). In Dedekind's cuts (1872), Cantor's 1891 diagonal argument, and Lawvere's 1969 fixed-point theorem [16], it acts as an exponential valuation apparatus over a continuous canvas ($\beth_1 = \Omega^{\beth_0}$, $\beth_2 = \Omega^{\beth_1}$).

How do these mutually incompatible operational regimes cohere into a single mathematical subject? Received philosophy of mathematics offers two equally flat answers:

1. **Ontological Monism (Set-Theoretic Platonism):** The token ∞ is treated as an ambiguous surface sign whose genuine referents reside in a single, pre-existing cumulative hierarchy $V = \bigcup_{\alpha \in \mathrm{On}} V_\alpha$. Concrete operational apparatuses (Hilbert's Hotel, binary bisection trees, Boustrophedon grids, diagonal games) are dismissed as heuristic scaffolding used to point toward static sub-levels of V ($\aleph_\alpha \leq \beth_\alpha$ inside $\mathrm{On} \subseteq V$). Yet this encoding achieves unity only by erasing operational structure: identifying the first infinite ordinal ω, the first infinite cardinal $\aleph_0$, and the countable valuation carrier $\beth_0$ as the exact same von Neumann set in V conceals the fact that the operations governing them obey equational laws that contradict each other once they are written with one sign.

2. **Syntactic Formalism and Deflationary Fragmentation:** At the opposite pole, formalism reduces mathematical sense to deductive derivability within a fixed axiomatic schema such as ZFC, while late-Wittgensteinian deflationism (*Remarks on the Foundations of Mathematics* [32]) treats each calculus as a self-contained "grammar" with no overarching mathematical referent. If Platonism collapses distinct operational worlds into a single flat ontology, deflationism shatters mathematical practice into unglued homonyms.

In the philosophy of natural language, Wittgenstein broke the deadlock between mentalism and logical atomism with the dictum that meaning is use within a rule-governed language-game [31] (compare Lakatos's analysis of informal proof dynamics [15] and Thurston's account of how mathematical understanding advances through shared mental models [29]). Mathematics, however, does not merely deploy words in social conversation; it invents explicit, self-contained operational miniatures—what we call **semantic toys**—with which mathematicians play, build, rearrange, and explore [21, 24, 27]. When Archimedes fills the universe with grains of sand, when Galileo pairs integers with their squares, when Hilbert shifts guests down an infinite corridor of hotel rooms, when Dedekind slices a stick into left and right halves, or when Cantor lines people up in a waiting queue and asks what happens when someone

steps behind the entire line, they are not merely illustrating pre-existing set-theoretic axioms. They are inhabiting the empty token ∞ with distinct **worlds of generative play**. We therefore formulate our foundational thesis:

> **The Principle of the Semantic Toy**. *In mathematics, meaning is the semantic toy.* A mathematical sign σ is an uninterpreted syntactic token until inhabited by a *semantic toy*: a constructive operational miniature defined by its material configurations ($\mathcal{X}_T$), its generative moves of play ($\mathcal{E}_T$), and its internal symmetries of rearrangement ($\mathcal{W}_T$). The local mathematical sense of σ inside a toy T is the localized meaning-fibre $\mathcal{M}(T) = \mathrm{Sk}(\mathrm{Pos}(\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]))$ of configurations modulo reversible play. A global mathematical concept is neither a single Platonic object nor a flat axiomatic pushout, but the **Grothendieck opfibration**
>
> $$p : \int_{\mathcal{T}_\infty} \mathcal{M} \longrightarrow \mathcal{T}_\infty$$
>
> gluing the local meaning-fibres of all semantic toys inhabiting σ across history and across operational stances.

1.2 Four Philosophical Interventions

Before developing the categorical formalism, we situate the Principle of the Semantic Toy relative to four central debates in the philosophy of mathematics:

1. Against Verificationism: Why a Semantic Toy Is a World of Generative Play, Not a Test. It is tempting for philosophers steeped in Dummettian anti-realism, logical positivism, or game-theoretic verificationism to confuse a *semantic toy* with a *verification test*—reading our account as the claim that mathematical objects are determined by "what challenges or tests a referee is permitted to run." That reading gets the philosophical priority backward.

A toy is not an examination; it is a **generative medium of play**. When a child plays with wooden blocks, or when a mathematician plays with Hilbert's Grand Hotel (T_{ticket}), a waiting line (T_{queue}), or a painted ruler (T_{ruler}), the toy is constituted first and foremost by **what pieces exist on the board** ($\mathcal{X}_T$), **what generative moves you can make to build new configurations** ($\mathcal{E}_T$), and **what counts as a natural, reversible rearrangement of the pieces** ($\mathcal{W}_T$). For example:

- In the **Ticket Toy (T_{ticket})**, the pieces are unordered collections of numbered tickets, the generative moves add new guests or lay tickets out in two-dimensional grids, and a reversible rearrangement is any one-to-one re-dealing of tickets.
- In the **Queue Toy (T_{queue})**, the pieces are well-ordered waiting lines in which *who stands ahead of whom* is part of the material piece itself; the generative moves place a newcomer at the very front of the line ($\mathbf{1} \oplus \alpha$) or behind the entire line ($\alpha \oplus \mathbf{1}$); and a reversible rearrangement is an order-preserving slide.
- In the **Ruler Toy (T_{ruler})**, the pieces are continuous rulers or canvases, and the generative move ($V \mapsto \Omega^V$) cuts or paints every point of the previous stage in two colours.

Why, then, do adversarial "challenges" (such as Cantor's diagonal flip or the pigeonhole obstruction on $\omega + 1$) appear in mathematics at all? Not because meaning is defined by testing, but because when we play with a toy and apply a generative move $E \in \mathcal{E}_T$ to a configuration x, one of two structural phenomena occurs:

(a) **Symmetry Absorption ($E(x) \sim_T x$)**: The newly built configuration $E(x)$ can be folded back into x by one of the toy's own reversible symmetries in $\mathcal{W}_T$. When a new guest arrives at Hilbert's Hotel ($\mathbb{N}^+ \sqcup \mathbf{1}$), shifting every guest one room down ($n \mapsto n+1$) is a lawful symmetry of ticket-play that absorbs the newcomer into the same hotel ($\aleph_0 + 1 = \aleph_0$). Similarly, when a newcomer steps to the *front* of an infinite queue ($\mathbf{1} \oplus \omega$), the order-preserving slide absorbs them ($1 + \omega = \omega$).

(b) **Generative Escape ($x <_T E(x)$)**: The generative move $E(x)$ builds a configuration so structurally rich that **no symmetry of play in $\mathcal{W}_T$ can fold it back into** x. When a newcomer steps

behind the entire infinite queue ($\omega \oplus \mathbf{1}$), they stand behind infinitely many people, whereas every position in ω stands behind only finitely many; no order-preserving slide can squeeze $\omega \oplus \mathbf{1}$ back into ω, so the move escapes to a genuinely new rung ($\omega + 1 > \omega$). Likewise, in the Ruler Toy, cutting or painting a domain ($V \mapsto \Omega^V$) escapes any discrete indexing by V ($\beth_0 < \beth_1 < \beth_2$).

Adversarial Proposer–Challenger games ($\mathcal{A}_T$; compare Lorenzen's dialogical logic [17] and Blass's game semantics [4]) are therefore **secondary diagnostics of generative escape**: they are how we feel the *structural friction* of the toy when a generative move has broken out of the old symmetry class and created a new scale of infinity.

2. Answering Deflationism via Grothendieck Gluing (On Wittgenstein's *RFM*). In *Remarks on the Foundations of Mathematics* (II, §§1–38) [32], Wittgenstein rightly observed that comparing $\aleph_0$ with $2^{\aleph_0}$ is not like comparing two finite piles of apples: the bijective ticket calculus and the diagonal power-set calculus operate by different rules. From this accurate operational observation, however, Wittgenstein drew a quietist, finitist conclusion: that Cantor's diagonal proof merely changes the subject and that transfinite set theory is a grammatical illusion.

Our framework accepts Wittgenstein's premise (different calculi are distinct semantic toys) while answering his deflationist conclusion. In modern category theory, when local mathematical worlds are governed by distinct internal rules of play, one does not discard them as illusions; one organizes them as fibres of a Grothendieck opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ [10, 2]. Intuitively, $\mathcal{T}_\infty$ is a floor plan of rooms (one room for each toy) connected by translation corridors, and $\mathcal{M}(T)$ is the vertical universe of mathematical objects generated by playing inside room T. Cantor's diagonal argument is not a grammatical sleight of hand; it is a rigorous cross-toy theorem demonstrating that when we carry the countable infinity $\aleph_0$ from the Ticket room into the Ruler room along the canonical corridor δ_{ruler} (where it becomes the discrete index carrier $\beth_0$), the Ruler Toy's generative valuation move $\beth_0 \mapsto \Omega^{\beth_0} = \beth_1$ escapes any indexing by $\beth_0$.

3. Inverting Benacerraf's Dilemma (Set-Theoretic Coding as Gauge-Fixing). Benacerraf's classic structuralist dilemma [3] observed that in finite arithmetic, a single operational structure $(\mathbb{N}, 0, S)$ admits many distinct set-theoretic encodings (such as Zermelo's $\{\{\emptyset\}\}$ versus von Neumann's $\{\emptyset, \{\emptyset\}\}$ for the number 2), so no single set can be "the" number 2.

At the transfinite level, the pathology is the exact inverse of Benacerraf's dilemma. Whereas finite arithmetic suffers from *one operational toy mapped to many set-theoretic codes*, transfinite set theory suffers from *many incompatible operational toys collapsed into a single set-theoretic code.* In ZFC, the Queue Toy's base configuration (queue, ω), the Ticket Toy's base configuration $(\mathsf{ticket}, \aleph_0)$, and the Ruler Toy's discrete index carrier $(\mathsf{ruler}, \beth_0)$ are all coded as the identical von Neumann ordinal $\omega \in V$. Set-theoretic coding acts as a flat projection (analogous to choosing a coordinate gauge in physics and mistaking the coordinates for the manifold) that obliterates the vertical fibre direction of $\int_{\mathcal{T}_\infty} \mathcal{M}$. Replacing flat set-theoretic coding with the opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ restores the distinct operational identities of ω, $\aleph_0$, and $\beth_0$ while recording their ground-level equivalence as a transport law along the corridors of $\mathcal{T}_\infty$.

4. Beyond the Ontological Multiverse vs. Universe Binary (Hamkins vs. Woodin). In contemporary philosophy of set theory, the independence of the Continuum Hypothesis ($\mathrm{CH} : \beth_1 = \aleph_1$) is predominantly debated in ontological terms: does there exist a single privileged Platonic universe V in which CH possesses a determinate truth value awaiting discovery via new large-cardinal or Ω-logic principles (Woodin [33]), or does set theory describe a pluralistic multiverse of equally real background universes $V[G]$ across which CH varies (Hamkins [11])?

The Semantic Toy framework offers a constructive, operational third way that requires neither Platonic realism about uncountable sets nor ontological pluralism about ghost universes. Both $\aleph_1$ and $\beth_1$ are generated by toys with explicit finite descriptions (the toys still run inside an ambient set theory:

Hartogs' construction uses Power Set, Separation and Replacement, and the valuation move uses Power Set):

- $\aleph_1$ is the first rung of the **Hartogs Escalator** on the Ticket Toy fibre $\mathcal{M}(\mathsf{ticket})$, which collects all countable well-ordered queue types modulo order-isomorphism;
- $\beth_1 = \Omega^{\beth_0}$ is the first rung of the **Exponential (Lawvere) Escalator** on the Ruler Toy fibre $\mathcal{M}(\mathsf{ruler})$, which collects binary valuations over the countable carrier.

Because T_{ticket} is an *extensional inventory toy* and T_{ruler} is an *intensional valuation toy*, they share a boundary condition at their countable ground floor ($\aleph_0 \simeq \beth_0$) but their escalators advance in orthogonal vertical fibres. The independence of CH from ZFC is the theorem of Gödel [9] and Cohen [5]; our framework does not derive it, it represents it. The base category $\mathcal{T}_\infty$ contains no corridor ruler $\to$ ticket pinning the exponential escalator to the Hartogs escalator, so the comparison has to be supplied from outside the toys, by a model. Inside each model U of ZFC the cardinalization $\theta_U : \mathcal{M}_U(\mathsf{ruler}) \to \mathcal{M}_U(\mathsf{ticket})$ exists, and there $\aleph_1 \leq \beth_1$ always holds; Gödel's constructible universe L and Cohen's forcing extensions are two *bridging meta-toys* in which $\theta_U(\beth_1)$ takes different values, constrained by Cantor's diagonal lower bound ($\beth_1 > \aleph_0$) and König's cofinality obstruction ($\mathrm{cf}(\beth_1) > \omega$).

2 Semantic Toys as Worlds of Generative Play, Symmetries of Action, and the Local Meaning-Fibre

2.1 Anatomy of a Semantic Toy: Pieces, Generative Moves, and Symmetries of Play

To make the Principle of the Semantic Toy accessible to both philosophers of mathematics and categorical logicians, let us unpack what happens when a mathematician constructs and plays with an operational miniature (such as shifting guests in Hilbert's Hotel, lining people up in a waiting queue, or bisecting a continuous ruler [27]). A semantic toy is built from three primary, constitutive layers, followed by a secondary diagnostic layer:

(a) **The Material Pieces and Board Configurations ($\mathcal{X}_T$):** Every toy begins by specifying what kind of concrete configurations live on its board and what structure those configurations carry. In Hilbert's Hotel (T_{ticket}), a configuration is an unordered collection of rooms/tickets; in the Queue Toy (T_{queue}), a configuration is a waiting line $(A, <_A)$ equipped with a strict precedence order; in the Ruler Toy (T_{ruler}), a configuration is a space of binary cuts or colourings Ω^V. Two distinct board states—such as "the full hotel $\mathbb{N}^+$" and "the full hotel plus one new guest in the lobby $\mathbb{N}^+ \sqcup \{\star\}$"—are two distinct raw configurations in $\mathcal{X}_T$. In Fregean terms, $\mathcal{X}_T$ is the realm of operational *modes of presentation* (*Sinne*).

(b) **The Generative Moves of Play ($\mathcal{E}_T$):** A toy is not a static museum display; it is defined by *what you can do with it.* Starting from the initial configuration $x_0(\sigma)$ that anchors the token σ inside the toy, the player applies constructive move generators $E_i : \mathcal{X}_T \to \mathcal{X}_T$ (together with colimits of chains of moves of any well-ordered length) to build new configurations: adjoining a new guest ($X \mapsto X \sqcup \mathbf{1}$), forming a two-dimensional grid of tickets ($X \mapsto X \times X$), placing a newcomer at the front or back of a queue ($\alpha \mapsto \mathbf{1} \oplus \alpha$ or $\alpha \mapsto \alpha \oplus \mathbf{1}$), collecting all well-ordered queues of a given ticket size into Hartogs' warehouse ($X \mapsto \aleph(X)$), or cutting/painting a domain in two colours ($V \mapsto \Omega^V$). The category $\mathcal{X}_T^{\langle\sigma\rangle}$ of configurations reachable from $x_0(\sigma)$ by playing these moves is the *generative horizon* of the toy.

(c) **The Internal Symmetries of Play ($\mathcal{W}_T$) and the Meaning-Fibre ($\mathcal{M}(T)$):** When do two raw configurations x and y built during play represent the *same* mathematical object (share the same local referent or *Bedeutung*) inside the toy T? Precisely when each can be absorbed into the other by a structure-preserving protocol ($x \preceq_T y$ and $y \preceq_T x$, written $x \sim_T y$). Notice that a generative extension morphism such as the canonical inclusion $\iota_1 : \mathbb{N}^+ \hookrightarrow \mathbb{N}^+ \sqcup \{\star\}$ in Hilbert's Hotel is *not* a surjective bijection in the raw protocol category $\mathcal{X}_{\mathsf{ticket}}$; rather, it is a **weak equivalence** ($w \in \mathcal{W}_{\mathsf{ticket}}$) because Hilbert's shift protocol $s : \mathbb{N}^+ \sqcup \{\star\} \hookrightarrow \mathbb{N}^+$ embeds the enlarged

configuration back into $\mathbb{N}^+$! To pass from the intensional category of raw configurations and individual protocols $\mathcal{X}_T^{\langle\sigma\rangle}$ (Fregean *Sinne*) to the invariant scale of local referents $\mathcal{M}(T)$ (Fregean *Bedeutungen*), we perform two canonical categorical steps:

(i) **Gabriel–Zisman Localization ($\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]$):** We formally invert all mutual-absorption morphisms $w \in \mathcal{W}_T$ [7], turning non-surjective generative extensions that admit a reverse absorption (such as $\iota_1 : \mathbb{N}^+ \hookrightarrow \mathbb{N}^+ \sqcup \{\star\}$) into formal isomorphisms; and

(ii) **Posetal Reflection and Skeleton ($\mathcal{M}(T) := \mathrm{Sk}(\mathrm{Pos}(\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]))$):** We 0-truncate parallel protocols between the same configurations into a single comparison relation ($\leq_T$) and pass to canonical isomorphism-class representatives $[x]_T$.

(d) **Symmetry Absorption vs. Generative Escape (The Diagnostic Game $\mathcal{A}_T$):** Once the pieces $\mathcal{X}_T$, the generative moves $\mathcal{E}_T$, and the weak equivalences $\mathcal{W}_T$ are in place, what happens when we apply a generative move $y = E(x)$? Either y can be folded back into x by a reverse protocol ($y \sim_T x$, **symmetry absorption**), or y genuinely outstrips x so that no lawful protocol can compress y back into x ($x <_T y$, **generative escape**). To witness whether y is absorbed by x or escapes x, we stage a two-player diagnostic game $\mathcal{A}_T(x, y)$ between a **Proposer (P)** (who attempts to fold y into x via a structure-preserving protocol f) and a **Challenger (C)** (who exhibits the specific piece of y that gets crushed, inverted, or left out by f, as in Gödel's Dialectica interpretation and Blass's game semantics [4]). Crucially, the game $\mathcal{A}_T$ does *not* create the toy; rather, $\mathcal{A}_T$ simply makes visible whether a generative move in $\mathcal{E}_T$ stays within the existing mutual-absorption class $\mathcal{W}_T$ or climbs to a strictly new rung of $\mathcal{M}(T)$.

Definition 2.1 (Semantic Toy Inhabiting a Token Sign). Let σ be a formal syntactic token (here $\sigma_\infty = \infty$). A *semantic toy T inhabiting σ* is a 5-tuple

$$T \;=\; \big(\mathcal{X}_T,\; x_0(\sigma),\; \mathcal{E}_T,\; \mathcal{W}_T,\; \mathcal{A}_T\big)$$

comprising the following constitutive and diagnostic data:

(i) **Configuration-and-Protocol Category ($\mathcal{X}_T$):** A category whose objects are concrete operational configurations (pointed state machines, unordered ticket sets, binary valuation spaces, or well-ordered waiting lines) and whose morphisms $f \in \mathcal{X}_T(x, y)$ are lawful structure-preserving protocols (pointed bisimulations in T_{loop}, injective ticket maps in T_{ticket}, zero-extended valuation embeddings in T_{ruler}, and strictly increasing maps in T_{queue}), closed under composition and identities.

(ii) **Anchor Configuration ($x_0(\sigma)$):** A distinguished initial configuration $x_0(\sigma) \in \mathrm{Ob}(\mathcal{X}_T)$ that inhabits the bare token σ at the ground entry of the toy.

(iii) **Generative Move Endofunctors ($\mathcal{E}_T$):** A family of constructive endofunctors $\mathcal{E}_T = \{E_i : \mathcal{X}_T \to \mathcal{X}_T\}_{i \in I}$ (equipped with extension morphisms $\iota_{E_i,x} : x \to E_i(x)$, natural where the move is functorial on raw configurations, and colimit cones for well-ordered chains of moves of any length (in the Queue Toy, chains of initial-segment embeddings); a move may instead be given only on the meaning-fibre, as an operation on classes $[x]_T$ with $[x]_T \leq_T E_i[x]_T$, which is how Hartogs' move enters the Ticket Toy, since on sets with all injections it has no natural unit $X \to \aleph(X)$: a non-trivial permutation of X acts trivially on $\aleph(X)$) representing the lawful moves of play by which new configurations are built. We write $\mathcal{X}_T^{\langle\sigma\rangle} \subseteq \mathcal{X}_T$ for the full subcategory of configurations generated from $x_0(\sigma)$ under finite and transfinite play of $\mathcal{E}_T$, and define the *absorption preorder* $x \preceq_T y \iff \mathcal{X}_T^{\langle\sigma\rangle}(x, y) \neq \emptyset$.

(iv) **Weak Equivalences / Mutual-Absorption Symmetries ($\mathcal{W}_T$):** We write $x \sim_T y$ whenever $x \preceq_T y$ and $y \preceq_T x$ (each configuration embeds into the other by a lawful protocol in $\mathcal{X}_T^{\langle\sigma\rangle}$). The wide subcategory of *weak equivalences* $\mathcal{W}_T \subseteq \mathcal{X}_T^{\langle\sigma\rangle}$ consists of all morphisms $w \in \mathcal{X}_T^{\langle\sigma\rangle}(x, y)$ between mutually absorbable configurations ($x \sim_T y$), i.e.,

$$\mathcal{W}_T(x, y) \;:=\; \big\{w \in \mathcal{X}_T^{\langle\sigma\rangle}(x, y) \;\big|\; \mathcal{X}_T^{\langle\sigma\rangle}(y, x) \neq \emptyset\big\}.$$

Because $\preceq_T$ is transitive, $\mathcal{W}_T$ is closed under composition and contains non-surjective generative inclusions $w : x \hookrightarrow E(x)$ whenever $E(x)$ can be folded back into x by a reverse protocol $g \in \mathcal{X}_T^{\langle\sigma\rangle}(E(x), x)$.

(v) **Diagnostic Obstruction Game ($\mathcal{A}_T$) for Generative Escape ($<_T$):** For any ordered pair $(x, y) \in \mathrm{Ob}(\mathcal{X}_T^{\langle\sigma\rangle})^2$, the diagnostic game $\mathcal{A}_T(x, y) = (M_{\mathsf{P}}(x, y), M_{\mathsf{C}}(x, y), u_{x,y})$ tests whether x suffices to absorb or cover y under the rules of T:

- A Proposer strategy $f \in M_{\mathsf{P}}(x, y)$ is a candidate protocol claiming to embed or index y inside x while respecting the toy's structure (a map $f : y \to x$ tested for injectivity/order-preservation in $\mathcal{A}_{\mathsf{queue}}(x, y)$ and $\mathcal{A}_{\mathsf{ticket}}(x, y)$, or a point-indexing catalogue $e : x \to y$ in $\mathcal{A}_{\mathsf{ruler}}(x, y)$).
- A Challenger counter-strategy $\chi \in M_{\mathsf{C}}(x, y)$ is an obstruction operator assigning to each candidate $f \in M_{\mathsf{P}}(x, y)$ a concrete structural defect $\chi(f)$ (a seat collision, an order inversion, or an omitted valuation) witnessing that f fails to absorb y into x.
- A winning Proposer strategy in $\mathcal{A}_T(x, y)$ yields a valid absorption protocol in $\mathcal{X}_T^{\langle\sigma\rangle}(y, x)$ ($y \preceq_T x$). Conversely, we write $x <_T y$ (*generative escape*) if $x \preceq_T y$ while in $\mathcal{A}_T(x, y)$ the Challenger C possesses a *uniform winning obstruction operator* $\chi_{x,y} \in M_{\mathsf{C}}(x, y)$ ($u_{x,y}(f, \chi_{x,y}) = \mathsf{C}$ for all $f \in M_{\mathsf{P}}(x, y)$), certifying $\mathcal{X}_T^{\langle\sigma\rangle}(y, x) = \emptyset$ ($y \not\preceq_T x$).

Definition 2.2 (Local Meaning-Fibre via Posetal Gabriel–Zisman Localization). Given a semantic toy T, let $\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]$ be the Gabriel–Zisman localization [7] of the protocol category $\mathcal{X}_T^{\langle\sigma\rangle}$ at its subcategory of weak equivalences $\mathcal{W}_T$. The *local meaning-fibre* $\mathcal{M}(T)$ of T is the skeletal posetal reflection of this localization:

$$\mathcal{M}(T) \;:=\; \mathrm{Sk}\Big(\mathrm{Pos}\big(\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]\big)\Big), \tag{1}$$

equipped with the canonical projection functor $L_T : \mathcal{X}_T^{\langle\sigma\rangle} \to \mathcal{M}(T)$. Here Sk denotes the poset of isomorphism classes $[x]_T$, not a chosen skeleton, so L_T involves no choice of representatives; by Lemma 2.3 it is the quotient of the reachability preorder $\preceq_T$ by $\sim_T$. By construction, $\mathcal{M}(T)$ is a thin skeletal category (a partially ordered class of local referents $[x]_T$) in which:

(a) Every weak equivalence $w \in \mathcal{W}_T(x, y)$ (including non-surjective generative extensions $x \hookrightarrow E(x)$ that admit a reverse absorption protocol $E(x) \hookrightarrow x$) is sent to the identity morphism $L_T(w) = \mathrm{id}_{[x]_T}$ on $[x]_T = [y]_T$ (the localization makes w an isomorphism; it becomes an identity after the skeletal posetal reflection);

(b) Parallel protocols $f, f' \in \mathcal{X}_T^{\langle\sigma\rangle}(x, y)$ are 0-truncated to the unique comparison arrow $[x]_T \leq_T [y]_T$.

2.2 Soundness of Strict Escape and Symmetry Relaxation Under Forgetful Translations

Definitions 2.1 and 2.2 resolve two fundamental questions at once: first, why adjoining formal inverses $\mathcal{W}_T^{-1}$ in the localization never accidentally collapses a strict generative escape $x <_T y$; and second, why stripping structure from the pieces of a toy (such as passing from the Queue Toy T_{queue} to the Ticket Toy T_{ticket}) relaxes its symmetries ($\mathcal{W}_{T_1} \subsetneq \mathcal{W}_{T_2}$) and folds strict ordinal inequalities ($\omega <_{\mathsf{queue}} \omega + 1$) into cardinal identities ($\aleph_0 = \aleph_0 + 1$).

Lemma 2.3 (Soundness of Localization and Forgetful Symmetry Relaxation). *(i)* ***Non-Collapse of Strict Escape ($x <_T y$):*** *For any semantic toy T and any configurations $x, y \in \mathrm{Ob}(\mathcal{X}_T^{\langle\sigma\rangle})$, there is a morphism from x to y in the Gabriel–Zisman localization $\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]$ if and only if $x \preceq_T y$. Consequently, two objects x, y become isomorphic in $\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]$ (and equal in $\mathcal{M}(T)$) if and only if $x \sim_T y$, and whenever $x <_T y$ (witnessed by a winning Challenger obstruction in $\mathcal{A}_T(x, y)$), the strict inequality $[x]_T <_T [y]_T$ is preserved in $\mathcal{M}(T)$.*

(ii) ***Forgetful Toy Translations:*** *Let $T_1 = (\mathcal{X}_1, x_0, \mathcal{E}_1, \mathcal{W}_1, \mathcal{A}_1)$ and $T_2 = (\mathcal{X}_2, y_0, \mathcal{E}_2, \mathcal{W}_2, \mathcal{A}_2)$ be two semantic toys linked by a structure-forgetting functor $F : \mathcal{X}_1^{\langle\sigma\rangle} \to \mathcal{X}_2^{\langle\sigma\rangle}$ (read game-*

theoretically, the Challenger in T_2 may object to fewer things). Then $F(\mathcal{W}_1) \subseteq \mathcal{W}_2$, and the universal property of localization and posetal reflection induces a canonical covariant functor between their meaning-fibres:

$$\Pi_{1\to 2} : \mathcal{M}(T_1) \longrightarrow \mathcal{M}(T_2), \qquad [x]_{T_1} \longmapsto [F(x)]_{T_2}.$$

Whenever a generative extension $\iota : x \hookrightarrow y$ achieves strict escape $x <_{T_1} y$ in T_1 ($\mathcal{X}_1(y, x) = \emptyset$), but $F(y)$ admits a reverse protocol $g \in \mathcal{X}_2(F(y), F(x))$ in T_2, we have $F(\iota) \in \mathcal{W}_2(F(x), F(y))$ and the strict inequality $[x]_{T_1} <_{T_1} [y]_{T_1}$ in $\mathcal{M}(T_1)$ maps to the identity $\Pi_{1\to 2}([x]_{T_1}) = \Pi_{1\to 2}([y]_{T_1})$ in $\mathcal{M}(T_2)$.

Proof. (i) Every morphism in $\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}](x, y)$ is represented by a finite zigzag of forward arrows $f_i \in \mathcal{X}_T^{\langle\sigma\rangle}(a_i, b_i)$ and backward weak equivalences w_j^{-1} with $w_j \in \mathcal{W}_T(c_j, d_j)$. For each forward arrow f_i, we have $a_i \preceq_T b_i$ by definition; for each backward arrow w_j^{-1}, since $w_j \in \mathcal{W}_T(c_j, d_j)$ implies $\mathcal{X}_T^{\langle\sigma\rangle}(d_j, c_j) \neq \emptyset$, we also have $d_j \preceq_T c_j$. By transitivity of $\preceq_T$ under composition in $\mathcal{X}_T^{\langle\sigma\rangle}$, any zigzag from x to y implies $x \preceq_T y$. Therefore, x and y are isomorphic in $\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]$ if and only if $x \preceq_T y$ and $y \preceq_T x$ ($x \sim_T y$). If $x <_T y$, then $y \not\preceq_T x$, so no zigzag from y to x exists in $\mathcal{X}_T^{\langle\sigma\rangle}[\mathcal{W}_T^{-1}]$, and $[x]_T <_T [y]_T$ in $\mathcal{M}(T)$.

(ii) If $w \in \mathcal{W}_1(x, y)$, there exists $h \in \mathcal{X}_1^{\langle\sigma\rangle}(y, x)$, so $F(h) \in \mathcal{X}_2^{\langle\sigma\rangle}(F(y), F(x))$ witnesses $F(w) \in \mathcal{W}_2(F(x), F(y))$. Hence $L_2 \circ F : \mathcal{X}_1^{\langle\sigma\rangle} \to \mathcal{M}(T_2)$ sends every morphism in $\mathcal{W}_1$ to an identity in the thin poset $\mathcal{M}(T_2)$, factoring uniquely through $\mathcal{M}(T_1)$ via $\Pi_{1\to 2}([x]_{T_1}) = [F(x)]_{T_2}$. If $x <_{T_1} y$ via $\iota : x \hookrightarrow y$ while $\mathcal{X}_2(F(y), F(x))$ is inhabited by a rearrangement g, then $F(\iota) \in \mathcal{W}_2$, so $\Pi_{1\to 2}([x]_{T_1}) = [F(x)]_{T_2} = [F(y)]_{T_2} = \Pi_{1\to 2}([y]_{T_1})$. □

Remark 2.4 (What Lemma 2.3 uses, and what the games add). Part (i) depends on the particular class $\mathcal{W}_T$ of Definition 2.1: a morphism is inverted only when its target already maps back to its source, so every backward step in a zigzag is matched by a forward protocol. It is not a general property of Gabriel–Zisman localization. The diagnostic games add no information to $\mathcal{M}(T)$: a uniform winning Challenger operator is a certificate that $\mathcal{X}_T^{\langle\sigma\rangle}(y, x)$ is empty. We keep the game layer $\mathcal{A}_T$ as an expository device, because it names the concrete defect (the collision, the inversion, the omitted valuation) that each classical proof exhibits.

3 The Genealogy of σ_∞ as Worlds of Constructive Play

Viewed through Definition 2.1 and Lemma 2.3, the history of the infinite is not a gradual discovery of higher floors in a static Platonic building, nor a sequence of arbitrary axiomatic conventions. It is a sequence of creative inventions in which mathematicians inhabited the token $\sigma_\infty = \infty$ with new **semantic toys**—new kinds of pieces ($\mathcal{X}_T$), new generative moves of play ($\mathcal{E}_T$), and new symmetries of rearrangement ($\mathcal{W}_T$)—each opening up a vertical world of meaning $\mathcal{M}(T)$ (Table 1).

3.1 Toy 0: From Finite Exhaustion (T_{fin}) to the Coinductive Loop Toy (T_{loop})

In Archimedes' *Sand Reckoner* and modern large-number hierarchies (10^6, 10^{100}, $10^{10^{100}}$), the pieces are finite tally counters bounded by physical inscription horizons (such as the $\approx 10^{80}$ baryons in the observable cosmos). The philosophical function of T_{fin} is negative: exhausting physical matter is not mathematical infinity, because every counter in T_{fin} eventually halts at 0 under repeated countdown steps $n \mapsto n - 1$.

The transfinite proper begins with the **Loop Toy** T_{loop}, corresponding to Aristotle's potential infinite (*Physics* III.6) and Wallis's lemniscate ∞ (*De sectionibus conicis*, 1655). To understand T_{loop} mathematically, it is helpful to contrast **algebras** (which build objects up from a base, like $0, S(0), S(S(0))$)

Table 1. Genealogy of semantic toys inhabiting the token $\sigma_\infty = \infty$, classified by material pieces $\mathcal{X}_T$, generative moves of play $\mathcal{E}_T$, internal symmetries $\mathcal{W}_T$ / escape obstructions $\mathcal{A}_T$, and localized objects in $\mathcal{M}(T)$.

Toy $T \in \mathcal{T}_\infty$	Historical Origin	Material Pieces $\mathcal{X}_T$	Generative Play $\mathcal{E}_T$	Symmetries $\mathcal{W}_T$ & Escape Friction $\mathcal{A}_T$	Invariants in $\mathcal{M}(T)$
T_{fin} (Exhaustion)	Archimedes (*Sand Reckoner*)	Bounded counters $N < B$	$n \mapsto n+1, b \mapsto 10^b$	Finite tallying; halts at physical atom bound B	$10^6 < 10^{100} < 10^{10^{100}}$
T_{loop} (Loop / Step)	Aristotle, Wallis (1655)	Non-halting coalgebras $(S, s_0, \text{next} : S \to S)$	Step continuation next^k	Pointed bisimilarity to the terminal loop $(\mathbf{1}, \star, \text{id})$	Single coalgebraic behavior σ_∞
T_{ticket} (Ticket Toy)	Galileo (1638), Cantor (1874), Hilbert (1924)	Unordered ticket bags / hotel rooms	Shift $X \sqcup \mathbf{1}$, grid $X \times X$, Hartogs $\aleph(X)$	Symmetry: mutual injection; Escape: Hartogs $\aleph(X)$	$\aleph_0 + 1 = \aleph_0, \lvert\mathbb{Q}\rvert = \aleph_0$, $\aleph_0 < \aleph_1 < \cdots < \aleph_\omega$
T_{ruler} (Ruler / Paint Toy)	Dedekind (1872), Cantor (1891), Lawvere (1969)	Valuation spaces $\mathbf{2}^V$ (cuts, 2-colourings)	Exponential valuation $V \mapsto \mathbf{2}^V$	Escape: Lawvere diagonal flip $\sigma \circ \widehat{e} \circ \Delta_V$	$\beth_0 < \beth_1 < \beth_2 < \ldots$, $\beth_1^{\aleph_0} = \beth_1$
T_{queue} (Queue Toy)	Cantor (1883), Hausdorff (1908), Gentzen (1936)	Well-ordered waiting lines $(A, <)$	Front step $\mathbf{1} \oplus \alpha$, post-limit $\alpha \oplus \mathbf{1}, \omega^\alpha$	Symmetry: mutual order-embedding; Escape: $\alpha \oplus \mathbf{1}$	$1 + \omega = \omega < \omega + 1 < \omega^2 < \cdots < \varepsilon_0$
T_{force} (Bridge Meta-Toy)	Gödel (L, 1940), Cohen ($\mathbf{Sh}(\mathbb{P})$, 1963), König (1905)	Topos / ZFC charts $U \in \mathcal{S}$	Poset extension $V \rightsquigarrow V[G]$	Cross-toy bridge $\theta_U(\beth_1) = \aleph_{\gamma(U)}$	$\gamma(L) = 1, \gamma(\mathbf{Sh}(\mathbb{P}_\beta)) = \beta$ over a GCH ground, $\text{cf}(\theta_U(\beth_1)) > \omega$

with **coalgebras** (which observe a dynamical process moving forward step by step). A general step-machine with possible termination is a pointed coalgebra $(S, s_0, \text{next} : S \to \mathbf{1} \sqcup S)$ for the maybe-functor $F(X) = \mathbf{1} \sqcup X$, whose terminal coalgebra $\nu X.\,(\mathbf{1} \sqcup X) \cong \mathbb{N} \sqcup \{\infty\}$ records finite halting step-counts $n \in \mathbb{N}$ together with the unique non-halting behavior ∞.

- **Pieces and Play ($\mathcal{X}_{\text{loop}}, \mathcal{E}_{\text{loop}}$):** The Loop Toy T_{loop} consists of *non-halting* pointed state machines $(S, s_0, \text{next} : S \to S)$—pointed coalgebras for the identity functor $\text{Id}(X) = X$—where playing the toy consists in pressing the step-generator next indefinitely. A protocol $R : (S, s_0, \text{next}) \to (S', s_0', \text{next}')$ is a *pointed bisimulation*: a relation $R \subseteq S \times S'$ with $(s_0, s_0') \in R$ and $(\text{next}(s), \text{next}'(s')) \in R$ whenever $(s, s') \in R$. Bisimulations compose by relational composition, and the diagonal is the identity.
- **Weak Equivalences ($\mathcal{W}_{\text{loop}}$):** Even when two non-halting machines are non-isomorphic as raw state spaces (for instance, the one-state self-loop $(\{\star\}, \star, \text{id})$ versus the unrolled counter $(\mathbb{N}, 0, n \mapsto n+1)$), both admit the unique homomorphism $! : (S, s_0, \text{next}) \to (\mathbf{1}, \star, \text{id})$ into the terminal coalgebra $\nu X.\, X \cong \mathbf{1}$ of the identity functor, whose graph is a pointed bisimulation. Its converse $\{(\star, s) \mid s \in S\}$ is a pointed bisimulation in the other direction. (A homomorphism back would not exist: $h : (\mathbf{1}, \star, \text{id}) \to (\mathbb{N}, 0, n \mapsto n+1)$ would need $h(\star) = h(\star) + 1$.) So every non-halting machine is mutually absorbable with the terminal loop $(\mathbf{1}, \star, \text{id})$, and every protocol between machines lies in $\mathcal{W}_{\text{loop}}$.

Consequently, the posetal localization $\mathcal{M}(T_{\text{loop}}) = \text{Sk}(\text{Pos}(\mathcal{X}_{\text{loop}}[\mathcal{W}_{\text{loop}}^{-1}]))$ contracts to a single object $\sigma_\infty = [\mathbf{1}]_{\text{loop}}$. This explains a striking historical fact: **as long as mathematics plays only with the Loop Toy of unending continuation, "infinity" has no plural**. Every forever-loop exhibits the exact same coalgebraic behavior.

3.2 Toy 1: The Ticket-Dispenser Toy (T_{ticket}) and Bijective Symmetries of Play

The seventeenth- and nineteenth-century breakthrough initiated by Galileo (1638), Bolzano, and Cantor (1874) and dramatized in Hilbert's Grand Hotel (1924) invents a new toy by unrolling the loop into an unordered roll of numbered tickets $x_0(\sigma_\infty) = \mathbb{N}^+$:

- **Pieces and Protocols ($\mathcal{X}_{\text{ticket}}$):** Objects are unordered collections of discrete tickets or hotel rooms, and morphisms in $\mathcal{X}_{\text{ticket}}(X, Y)$ are collision-free (injective) ticket assignments $f : X \hookrightarrow Y$.
- **Generative Play ($\mathcal{E}_{\text{ticket}}$):** Adjoining new guests ($X \mapsto X \sqcup \mathbf{1}$, $X \mapsto X \sqcup X$), laying tickets out in two-dimensional grids ($X \mapsto X \times X$), and collecting all well-ordered queues over X into Hartogs' warehouse ($X \mapsto \aleph(X)$).

- **Weak Equivalences ($\mathcal{W}_{\text{ticket}}$)**: An injective protocol $w : X \hookrightarrow Y$ belongs to $\mathcal{W}_{\text{ticket}}$ whenever there exists a reverse injective protocol $g : Y \hookrightarrow X$ (so $|X| = |Y|$ by the Cantor–Bernstein theorem).

Under posetal localization $\mathcal{M}(\text{ticket}) = \text{Sk}(\text{Pos}(\mathcal{X}_{\text{ticket}}^{\langle\sigma\rangle}[\mathcal{W}_{\text{ticket}}^{-1}]))$, any generative move $X \hookrightarrow E(X)$ that admits a reverse injection $E(X) \hookrightarrow X$ is *absorbed by symmetry*:

(a) **Hilbert Symmetry Absorption ($\aleph_0 + 1 = \aleph_0$, $\aleph_0 + \aleph_0 = \aleph_0$)**: When a new guest arrives ($Y = \mathbb{N}^+ \sqcup \{\star\}$), the canonical inclusion $\iota_1 : \mathbb{N}^+ \hookrightarrow \mathbb{N}^+ \sqcup \{\star\}$ is not surjective, but the Hilbert shift protocol $f(\star) = 1$, $f(n) = n + 1$ embeds Y back into $\mathbb{N}^+$ without collision; when a second infinite bus arrives ($Y = \mathbb{N}^+ \sqcup \mathbb{N}^+$), the odd/even interleave $f(1, n) = 2n$, $f(2, k) = 2k - 1$ embeds Y into $\mathbb{N}^+$. Thus $\iota_1 \in \mathcal{W}_{\text{ticket}}$, and localization yields $[\mathbb{N}^+ \sqcup \mathbf{1}]_{\text{ticket}} = [\mathbb{N}^+ \sqcup \mathbb{N}^+]_{\text{ticket}} = \aleph_0$ in $\mathcal{M}(\text{ticket})$.

(b) **Boustrophedon Grid Absorption ($|\mathbb{Q}| = \aleph_0$)**: Although the positive fractions $\mathbb{Q}_+$ appear infinitely denser than the integers on a geometric ruler, inside the Ticket Toy we can lay reduced ratios a/b ($\gcd(a, b) = 1$) on the two-dimensional grid $\mathbb{N}^+ \times \mathbb{N}^+$ and deal consecutive integer tickets along anti-diagonals $a + b = s$ while skipping non-coprime duplicates ($\gcd(a, b) > 1$):

$$\text{ticket}(a, b) \;=\; \text{next consecutive integer index} \quad \text{iff} \quad \gcd(a, b) = 1. \tag{2}$$

This skip-and-renumber snake-path defines an explicit bijection in $\mathcal{W}_{\text{ticket}}(\mathbb{N}^+, \mathbb{Q}_+)$, folding the rational grid back into $\aleph_0$. By contrast, Hartogs' warehouse move $X \mapsto \aleph(X)$ (Section 5) achieves *generative escape* ($\aleph_0 <_{\text{ticket}} \aleph_1 <_{\text{ticket}} \aleph_2 <_{\text{ticket}} \dots$).

3.3 Toy 2: The Ruler-Cutter and Canvas-Painter Toy (T_{ruler})

The second transfinite toy (Dedekind 1872, Cantor 1891, Lawvere 1969) replaces discrete ticket-dealing with a continuous geometric and combinatorial medium:

- **Pieces and Covariant Protocols ($\mathcal{X}_{\text{ruler}}$)**: Objects are valuation carriers V and spaces of binary cuts or colourings Ω^V, where $\Omega = \mathbf{2} = \{L, R\} \cong \{0, 1\}$. While exponential precomposition $q \mapsto q \circ f$ is contravariant on arbitrary maps, in $\mathcal{X}_{\text{ruler}}$ morphisms are *injective carrier embeddings* $i : A \hookrightarrow B$, which push binary valuations forward **covariantly** via zero-extension (direct-image valuation extension) $i_! : \mathbf{2}^A \hookrightarrow \mathbf{2}^B$, defined by $(i_!(q))(b) = q(a)$ if $b = i(a)$ and 0 otherwise.
- **Generative Play ($\mathcal{E}_{\text{ruler}}$)**: The **covariant exponential valuation endofunctor** $V \mapsto \mathbf{2}^V$ (equipped with the singleton-indicator embedding $\{\cdot\}_V : V \hookrightarrow \mathbf{2}^V$, $v \mapsto \delta_v$, and acting on injective embeddings $i : A \hookrightarrow B$ by $i_! : \mathbf{2}^A \hookrightarrow \mathbf{2}^B$), which takes an existing carrier V and forms the space of all ways of cutting or painting V.
- **Always-Escaping Escalation ($\mathcal{A}_{\text{ruler}}$)**: Unlike adding a ticket ($X \mapsto X \sqcup \mathbf{1}$), which is absorbed by Hilbert's shift, applying the valuation move $V \mapsto \mathbf{2}^V$ *always escapes* the previous stage ($V <_{\text{ruler}} \mathbf{2}^V$). The singleton embedding $\{\cdot\}_V : V \hookrightarrow \mathbf{2}^V$ ensures $V \preceq_{\text{ruler}} \mathbf{2}^V$, while whenever a Proposer tries to cover $\mathbf{2}^V$ using a catalogue $e : V \to \mathbf{2}^V$ indexed by the old piece V, the geometry of the toy induces the diagonal obstruction $\chi(e) \in \mathbf{2}^V$ that flips the value assigned to each $v \in V$ at locus v itself ($\forall v \in V,\ \chi(e) \neq e(v)$).

Playing the move generator $V \mapsto \mathbf{2}^V$ repeatedly from the discrete carrier $V_0 = \beth_0$ builds the beth tower:

- **Stage 1 ($\beth_1 = \mathbf{2}^{\beth_0}$, Binary Ruler Cuts)**: Playing $V_0 \mapsto \mathbf{2}^{V_0}$ builds the Cantor space $\{L, R\}^\omega$ of infinite binary bisection streams (which maps onto the geometric interval $[0, 1]$ modulo the countable set of dyadic rationals having two expansions, so $|[0, 1]| = |\mathbf{2}^{\beth_0}| = \beth_1$). Any countable list of binary streams $r_1, r_2, \dots \in \{L, R\}^\omega$ misses the diagonal stream $\chi(r)[i] = \text{flip}(r_i[i])$ (and on $[0, 1]$, avoiding $\{0, 1\}$-tail ambiguities via ternary/non-dyadic digits produces a point in $[0, 1]$ outside the list). Even if the player borrows Hilbert's shift move to prepend $\chi(r)$ at the top of the list ($r' = (\chi(r), r_1, r_2, \dots)$), the diagonal operator immediately generates a fresh omitted stream $\chi(r')$, witnessing $\beth_0 <_{\text{ruler}} \beth_1$.
- **Stage 2 ($\beth_2 = \mathbf{2}^{\beth_1}$, Two-Colourings of the Continuum)**: Playing the move a second time ($V_1 \mapsto \mathbf{2}^{V_1}$) paints the entire continuum V_1 in two colours. Any catalogue $\{C_x\}_{x \in V_1}$ of paintings indexed

by points of V_1 misses the diagonal painting

$$\chi(C)(x) \;=\; 1 - C_x(x) \qquad \text{for all } x \in V_1, \tag{3}$$

which differs from painting C_x at point x itself, witnessing $\beth_1 <_{\mathsf{ruler}} \beth_2$.

3.4 Toy 3: The Queue-Jumping Toy (T_{queue}) and Order-Preserving Play

The third transfinite toy (Cantor 1883, Hausdorff 1908, Gentzen 1936) starts from a countable collection of positions $x_0(\sigma_\infty) = (\omega, <)$, but builds **waiting-line precedence** directly into the material pieces of the toy:

- **Pieces and Protocols ($\mathcal{X}_{\mathsf{queue}}$):** Objects are well-ordered waiting lines $(A, <_A)$, and morphisms $f \in \mathcal{X}_{\mathsf{queue}}(A, B)$ are strictly order-preserving embeddings ($a_1 <_A a_2 \implies f(a_1) <_B f(a_2)$).
- **Generative Play ($\mathcal{E}_{\mathsf{queue}}$):** Stepping to the very front of a queue ($\alpha \mapsto \mathbf{1} \oplus \alpha$), stepping behind the entire completed queue ($\alpha \mapsto \alpha \oplus \mathbf{1}$), concatenating queues ($\alpha \oplus \beta$), forming exponential towers ($\alpha \mapsto \omega^\alpha$), and taking unions of chains of initial segments of any well-ordered length. Transfinite play of successor and unions alone reaches every ordinal $\alpha \geq \omega$ of the fixed universe, so the generative horizon of ω is the class of all infinite ordinals there; likewise, in the Ticket Toy, Hartogs' move and suprema reach every $\aleph_\alpha$.
- **Weak Equivalences ($\mathcal{W}_{\mathsf{queue}}$):** Because the waiting-line order is part of the piece itself, a protocol $w : A \hookrightarrow B$ belongs to $\mathcal{W}_{\mathsf{queue}}$ if and only if B also order-embeds into A ($A \sim_{\mathsf{queue}} B$, which for well-orders holds if and only if A and B are order-isomorphic, $\mathrm{otp}(A) = \mathrm{otp}(B)$); jumping the line breaks the piece.

By Lemma 2.3, because the forgetful functor $F : \mathcal{X}_{\mathsf{queue}} \to \mathcal{X}_{\mathsf{ticket}}$ relaxes order-embeddings to arbitrary injections ($F(\mathcal{W}_{\mathsf{queue}}) \subsetneq \mathcal{W}_{\mathsf{ticket}}$), the posetal localization $\mathcal{M}(\mathsf{queue}) = \mathrm{Sk}(\mathrm{Pos}(\mathcal{X}_{\mathsf{queue}}^{\langle\sigma\rangle}[\mathcal{W}_{\mathsf{queue}}^{-1}])) = \mathbf{Ord}_{\leq}$ distinguishes between front-stepping (which is absorbed by symmetry) and back-stepping (which escapes to a new rung):

(a) **Front-Step Symmetry Absorption ($1 + \omega = \omega$):** When a newcomer $\star$ steps to the very front of the queue ($\mathbf{1} \oplus \omega$), the slide $f(\star) = 0$, $f(n) = n + 1$ keeps everyone in their exact relative order ($\star < 0 < 1 < \cdots \implies 0 < 1 < 2 < \dots$). Thus $f \in \mathcal{W}_{\mathsf{queue}}$ is a reversible symmetry of the queue, and $\mathbf{1} \oplus \omega = \omega$ in $\mathcal{M}(\mathsf{queue})$.

(b) **Post-Limit Generative Escape ($\omega + 1 > \omega$):** When a newcomer $\top$ steps *behind* the entire infinite queue ($\omega \oplus \mathbf{1}$), $\top$ has infinitely many people standing ahead of them, whereas every position $m \in \omega$ has only finitely many positions $\{0, 1, \dots, m-1\}$ ahead of it. If a Proposer tries to squeeze $\omega \oplus \mathbf{1}$ back into ω by placing $\top$ at some finite spot $m = f(\top) \in \omega$, the pigeonhole principle forces some person $n < \top$ from the infinite line to be pushed behind m ($f(n) > m = f(\top)$), breaking the waiting-line order. Thus $\omega \oplus \mathbf{1}$ escapes the symmetry class of ω, establishing $\omega <_{\mathsf{queue}} \omega + 1$.

4 The Cost of a Shared Signature, and the Grothendieck Opfibration

4.1 One Signature for Two Toys

We first record what goes wrong when the local truths of the Queue and Ticket Toys of Section 3 are merged over one shared signature, in which the anchor σ_∞ and the successor $- \oplus \mathbf{1}$ are each a single symbol. The point is where the collapse comes from.

Theorem 4.1 (Collapse under a Shared Signature). *Let $\Sigma = (\sigma_\infty, - \oplus \mathbf{1}, <)$ be the shared signature of an anchor, right extension by one new element (a unary symbol, written $x \oplus \mathbf{1}$; the finite configuration $\mathbf{1}$ is not itself an element of the fibres), and strict comparison. Let $\mathsf{Th}(T_{\mathsf{queue}})$ and $\mathsf{Th}(T_{\mathsf{ticket}})$ denote the Σ-theories valid in the localized meaning-fibres $\mathcal{M}(\mathsf{queue})$ and $\mathcal{M}(\mathsf{ticket})$ under the anchor interpretations $\sigma_\infty \mapsto \omega$ (with $\alpha \oplus \mathbf{1} = \alpha + 1$) and $\sigma_\infty \mapsto \aleph_0$ (with $\kappa \oplus \mathbf{1} = \kappa + 1$), respectively. Then:*

(i) The union $\mathsf{Th}_{\mathsf{flat}} = \mathsf{Th}(T_{\mathsf{queue}}) \cup \mathsf{Th}(T_{\mathsf{ticket}})$ of the two theories over the shared signature Σ has no model in which $<$ is irreflexive.

(ii) Extend Σ by an ω-ary supremum symbol $\sup_n$*, and let the* countable ordinal terms *(well-founded trees with countable branching, so an infinitary syntax) be generated from σ_∞ by $t \mapsto t \oplus \mathbf{1}$ and $(t_n)_{n<\omega} \mapsto \sup_n t_n$. Every countable ordinal $\alpha \geq \omega$ is the value in* **Ord** *of some such term (as $\alpha = \beta + 1$ or as the supremum of a cofinal ω-sequence); $\omega + 1$, $\omega \cdot 2$, ω^2, ω^ω and ε_0 are examples. In any Σ-structure satisfying the Ticket Toy's absorption law $\sigma_\infty \oplus \mathbf{1} = \sigma_\infty$ and the law that the supremum of a constant sequence is that constant, every countable ordinal term denotes the same element as σ_∞.*

Proof. (i) In $\mathcal{M}(\text{queue})$, post-limit generative escape on $\omega \oplus \mathbf{1}$ validates the sentence $\sigma_\infty < \sigma_\infty \oplus \mathbf{1}$. In $\mathcal{M}(\text{ticket})$, Hilbert shift symmetry absorption validates the equation $\sigma_\infty \oplus \mathbf{1} = \sigma_\infty$. In any Σ-structure $\mathfrak{A}$ modeling $\mathsf{Th}_{\text{flat}}$, congruence of $<$ with respect to $=$ yields $\sigma_\infty^{\mathfrak{A}} < \sigma_\infty^{\mathfrak{A}}$, violating the irreflexivity axiom $\forall x\, \neg(x < x)$. (ii) By induction on terms. The base case is σ_∞ itself. If t denotes σ_∞, then $t \oplus \mathbf{1}$ denotes $\sigma_\infty \oplus \mathbf{1} = \sigma_\infty$. If every t_n denotes σ_∞, then $\sup_n t_n$ is the supremum of a constant sequence and denotes σ_∞. □

Remark 4.2 (External Operation Tagging vs. Fibred Typing). Theorem 4.1 is about one shared signature, not about ZFC. ZFC codes $\omega = \aleph_0 = \beth_0$ as the same von Neumann set and avoids the contradiction by keeping the operations distinct ($+_{\text{ord}}$ versus $+_{\text{card}}$). The carrier alone then does not say which regime is in force; the operation applied to it does. Indexing the meaning of σ_∞ over the base category of toys $\mathcal{T}_\infty$ assigns each operational regime its own *typed meaning-fibre* $\mathcal{M}(T)$, so that (queue, ω), $(\text{ticket}, \aleph_0)$, and $(\text{ruler}, \beth_0)$ are distinct typed objects connected by explicit functorial transport.

4.2 The Base Category $\mathcal{T}_\infty$ and the Queue–Ticket Adjunction

Definition 4.3 (Base Category of Transfinite Semantic Toys $\mathcal{T}_\infty$). The *base category of transfinite semantic toys* $\mathcal{T}_\infty$ has three core transfinite objects queue (T_{queue}), ticket (T_{ticket}), and ruler (T_{ruler}) (entered from the coalgebraic root loop via the unrolling morphism $u_0 : \text{loop} \to \text{queue}$), with generating morphisms:

$$\text{queue} \underset{\iota_{\text{init}}}{\overset{\pi_{\text{ticket}}}{\rightleftarrows}} \text{ticket} \xrightarrow{\delta_{\text{ruler}}} \text{ruler}$$

subject to the relation $\pi_{\text{ticket}} \circ \iota_{\text{init}} = \text{id}_{\text{ticket}}$, where:

- $\pi_{\text{ticket}} : \text{queue} \to \text{ticket}$ is the *order-forgetting symmetry relaxation* (Lemma 2.3), which strips waiting-line precedence from the pieces and relaxes order-embeddings to arbitrary injections;
- $\iota_{\text{init}} : \text{ticket} \to \text{queue}$ is the *initial-ordinal realization*, which sends each cardinal scale to its minimal well-ordered representative; and
- $\delta_{\text{ruler}} : \text{ticket} \to \text{ruler}$ is the *discrete carrier inclusion*, which imports a cardinal scale as a discrete carrier in the valuation fibre.

Let us explain in plain mathematical and philosophical terms how the localized Queue Toy fibre $\mathcal{M}(\text{queue}) = \mathbf{Ord}_{\leq}$ (the thin skeletal category of infinite ordinals $\alpha \geq \omega$ ordered by $\alpha \leq \beta$) and the localized Ticket Toy fibre $\mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$ (the thin skeletal category of infinite well-ordered cardinals $\aleph_\gamma$ ordered by $\aleph_\gamma \leq \aleph_\delta$) interact across the pair of corridors $(\iota_{\text{init}}, \pi_{\text{ticket}})$.

First, observe why the two-step passage in Definition 2.2—from the raw protocol categories ($\mathcal{X}_{\text{queue}} = \mathbf{Ord}_{\text{emb}}$, $\mathcal{X}_{\text{ticket}} = \mathbf{Card}_{\text{inj}}$) to the thin posetal meaning-fibres ($\mathbf{Ord}_{\leq}$, $\mathbf{Card}_{\leq}$)—is mathematically essential. In the raw protocol categories, hom-sets record individual embedding functions, which have incompatible automorphism groups: for instance, on a two-element configuration, $|\mathbf{Ord}_{\text{emb}}(2,2)| = 1$ whereas $|\mathbf{Card}_{\text{inj}}(2,2)| = 2$, and on ω, $\mathbf{Card}_{\text{inj}}(\omega,\omega)$ contains non-identity permutations whereas every well-order has only the identity order-automorphism. Thus no functor $I : \mathbf{Card}_{\text{inj}} \to \mathbf{Ord}_{\text{emb}}$ satisfies $U \circ I \cong \text{id}$: a non-identity permutation σ of ω must go to an order-automorphism, hence to the identity, and naturality of $U \circ I \cong \text{id}$ would then force $\sigma = \text{id}$. Passing via posetal Gabriel–Zisman localization (Definition 2.2) to the thin meaning-fibres $\mathcal{M}(\text{queue}) = \mathbf{Ord}_{\leq}$ and $\mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$ 0-truncates parallel protocols to their invariant order relation ($\leq$), where the initial-ordinal realization and order-forgetting projection form an exact Galois adjunction:

When we apply the order-forgetting functor $\mathcal{M}(\pi_{\text{ticket}}) : \mathbf{Ord}_{\leq} \to \mathbf{Card}_{\leq}$, an entire infinite vertical column of distinct queues $(\omega, \omega+1, \omega\cdot 2, \omega^2, \ldots, \varepsilon_0)$ gets compressed onto the single ticket count $\aleph_0$. Conversely, given a ticket size $\aleph_\gamma \in \mathbf{Card}_{\leq}$, $\mathcal{M}(\iota_{\text{init}})(\aleph_\gamma) = \omega_\gamma$ selects the **initial ordinal** of size $\aleph_\gamma$: the shortest well-ordered queue with $\aleph_\gamma$ tickets. Because ω_γ sits at the very bottom of the vertical column of queues of size $\aleph_\gamma$, we have $\omega_\gamma \leq \alpha$ in $\mathbf{Ord}_{\leq}$ if and only if $\aleph_\gamma \leq |\alpha|$ in $\mathbf{Card}_{\leq}$, meaning that $\mathcal{M}(\iota_{\text{init}})$ **is left adjoint to** $\mathcal{M}(\pi_{\text{ticket}})$.

Proposition 4.4 (The Initial-Ordinal Galois Adjunction $\mathcal{M}(\iota_{\text{init}}) \dashv \mathcal{M}(\pi_{\text{ticket}})$). *On the thin skeletal meaning-fibres $\mathcal{M}(\text{queue}) = \mathbf{Ord}_{\leq}$ and $\mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$, let $\mathcal{M}(\pi_{\text{ticket}}) : \mathbf{Ord}_{\leq} \to \mathbf{Card}_{\leq}$ be the order-forgetting functor $\alpha \mapsto |\alpha|$, and let $\mathcal{M}(\iota_{\text{init}}) : \mathbf{Card}_{\leq} \hookrightarrow \mathbf{Ord}_{\leq}$ map each cardinal $\kappa = \aleph_\gamma$ to its initial ordinal $\omega_\gamma = \mu\alpha.\,(|\alpha| = \kappa)$ (the subscript indexes initial ordinals; it is not ordinal exponentiation ω^γ). Then $\mathcal{M}(\pi_{\text{ticket}}) \circ \mathcal{M}(\iota_{\text{init}}) = \mathrm{id}_{\mathbf{Card}_{\leq}}$ holds strictly, and $\mathcal{M}(\iota_{\text{init}})$ is left adjoint to $\mathcal{M}(\pi_{\text{ticket}})$:*

$$\mathcal{M}(\iota_{\text{init}}) \;\dashv\; \mathcal{M}(\pi_{\text{ticket}}) \;:\; \mathbf{Ord}_{\leq} \longrightarrow \mathbf{Card}_{\leq}. \tag{4}$$

That is, for every cardinal $\kappa \in \mathbf{Card}_{\leq}$ and every ordinal $\alpha \in \mathbf{Ord}_{\leq}$, there is a natural bijection of singleton/empty hom-sets (a Galois connection of posets):

$$\mathrm{Hom}_{\mathbf{Ord}_{\leq}}\big(\mathcal{M}(\iota_{\text{init}})(\kappa),\, \alpha\big) \;\cong\; \mathrm{Hom}_{\mathbf{Card}_{\leq}}\big(\kappa,\, \mathcal{M}(\pi_{\text{ticket}})(\alpha)\big), \qquad \omega_\gamma \leq \alpha \iff \aleph_\gamma \leq |\alpha|.$$

Furthermore, the unit of the adjunction $\eta_\kappa : \kappa \xrightarrow{=} \mathcal{M}(\pi_{\text{ticket}})(\mathcal{M}(\iota_{\text{init}})(\kappa))$ is the identity morphism on κ, while the counit $\epsilon_\alpha : \mathcal{M}(\iota_{\text{init}})(\mathcal{M}(\pi_{\text{ticket}})(\alpha)) \leq \alpha$ is the unique initial-segment inequality $\omega_\gamma \leq \alpha$ pointing from the floor of the vertical column (ω_γ) up to α for every ordinal α of cardinality $\aleph_\gamma$.

Proof. Because $\mathbf{Ord}_{\leq}$ and $\mathbf{Card}_{\leq}$ are thin skeletal poset categories, each hom-set contains at most one arrow. By definition of the initial ordinal $\omega_\gamma = \mathcal{M}(\iota_{\text{init}})(\aleph_\gamma)$, ω_γ is the least ordinal of cardinality $\aleph_\gamma$, so $|\omega_\gamma| = \aleph_\gamma$ (giving $\mathcal{M}(\pi_{\text{ticket}}) \circ \mathcal{M}(\iota_{\text{init}}) = \mathrm{id}_{\mathbf{Card}_{\leq}}$ on objects and arrows) and $\omega_\gamma \leq \alpha$ in $\mathbf{Ord}_{\leq}$ if and only if $\aleph_\gamma = |\omega_\gamma| \leq |\alpha| = \mathcal{M}(\pi_{\text{ticket}})(\alpha)$ in $\mathbf{Card}_{\leq}$. Since both hom-sets have size 1 when the inequality holds and size 0 otherwise, this gives the natural hom-set bijection $\mathrm{Hom}_{\mathbf{Ord}_{\leq}}(\omega_\gamma, \alpha) \cong \mathrm{Hom}_{\mathbf{Card}_{\leq}}(\aleph_\gamma, |\alpha|)$. For any α with $|\alpha| = \aleph_\gamma$, the counit $\epsilon_\alpha : \omega_\gamma \leq \alpha$ is the unique comparison arrow in $\mathbf{Ord}_{\leq}$, which is an identity in $\mathbf{Ord}_{\leq}$ if and only if $\alpha = \omega_\gamma$. □

4.3 The Grothendieck Opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$

Let us now unpack the central categorical construction of this paper—the **Grothendieck opfibration** $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$—so that its geometric and philosophical meaning is clear:

- **What is the Grothendieck construction ($\int_{\mathcal{T}_\infty} \mathcal{M}$)?** Imagine an architectural building whose ground-floor rooms are the semantic toys $T \in \mathcal{T}_\infty$ (queue, ticket, ruler). Above each room T rises a vertical tower of mathematical configurations: its local meaning-fibre $\mathcal{M}(T)$. The Grothendieck construction $\int_{\mathcal{T}_\infty} \mathcal{M}$ is simply the **entire multi-room building taken as a single total category**. Every object in $\int_{\mathcal{T}_\infty} \mathcal{M}$ carries a two-part address (T, x): first, the name of the room T it lives in; second, the configuration $x \in \mathcal{M}(T)$ inside that room. Because $(\text{queue}, \omega+1)$ lives in the queue room and $(\text{ticket}, \aleph_0)$ lives in the ticket room, they never collide or contradict each other.
- **What is an Opfibration, and why does $\mathcal{M}(\pi_{\text{ticket}})$ have no right adjoint?** When we walk along a translation corridor $u : T \to T'$ in the base floor plan (for instance, walking along $\pi_{\text{ticket}} : \text{queue} \to \text{ticket}$ by forgetting the queue order), our functor $\mathcal{M}(u) = u_! : \mathcal{M}(T) \to \mathcal{M}(T')$ carries objects *forward* in the same direction as the arrow u (sending $\omega+1 \mapsto \aleph_0$). In category theory, when a family of fibres is equipped with canonical forward transport along base arrows, the projection $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ is called a **Grothendieck opfibration** [10, 2]. Notice an illuminating asymmetry in the $\{\text{queue}, \text{ticket}\}$ sector: along $\iota_{\text{init}} : \text{ticket} \to \text{queue}$, the forward functor $\mathcal{M}(\iota_{\text{init}})$ has a right adjoint $\mathcal{M}(\pi_{\text{ticket}})$ (Proposition 4.4), so ι_{init} admits both cocartesian and cartesian lifts. Along $\pi_{\text{ticket}} : \text{queue} \to \text{ticket}$, however, the forward functor $\mathcal{M}(\pi_{\text{ticket}}) : \mathbf{Ord}_{\leq} \to \mathbf{Card}_{\leq}$ has a *left* adjoint $\mathcal{M}(\iota_{\text{init}})$ (selecting the *minimum* ordinal ω_γ of cardinality $\aleph_\gamma$) but **no right adjoint**!

Why? Because a right adjoint $R(\aleph_\gamma)$ would have to be the *greatest* ordinal of cardinality $\leq \aleph_\gamma$ ($|\alpha| \leq \aleph_\gamma \iff \alpha \leq R(\aleph_\gamma)$), and the vertical fibre $[\omega, \omega_{\gamma+1})$ has no maximum element—its supremum escapes to the Hartogs successor $\omega_{\gamma+1}$ (Proposition 5.2)!

- **What is a "Cocartesian Lift" ($\widetilde{u}_x$)?** Given an object (T, x) in room T and a corridor $u : T \to T'$ to room T', the arrow $\widetilde{u}_x : (T, x) \to (T', u_!(x))$ in the total building $\int_{\mathcal{T}_\infty} \mathcal{M}$ is called the **cocartesian lift** of u at x. Philosophically, it represents *pure translation without extra modification*: it carries x across the threshold from Toy T into Toy T' and lands on the exact image $u_!(x)$ dictated by the translation rule u. Any other cross-room morphism from (T, x) to some (T', y) factors uniquely as first crossing the threshold via the cocartesian lift $\widetilde{u}_x$ to $(T', u_!(x))$, and then performing an internal proof $\varphi : u_!(x) \leq_{T'} y$ vertically inside room T'.

Definition 4.5 (The Grothendieck Opfibration of Semantic Toys). Fix a Grothendieck universe $\mathcal{U}$ bounding size: the fibres are the ordinals, cardinals and valuation carriers of $\mathcal{U}$, and the "any length" of play means any length in $\mathcal{U}$. A universe is closed under power set and replacement, so Hartogs' move and $V \mapsto \mathbf{2}^V$ stay inside it, and the fibres are small relative to a larger universe, as Thomason's theorem requires. Let $\mathcal{M} : \mathcal{T}_\infty \to \mathbf{Cat}$ be the strict covariant functor defined on objects and generating morphisms by:

- $\mathcal{M}(\mathsf{loop}) = \{\sigma_\infty\}$ (the terminal one-object category), $\mathcal{M}(\mathsf{queue}) = \mathbf{Ord}_\leq$ (infinite ordinals $\alpha \geq \omega$ ordered by $\leq$; the generative horizon of ω under successor and unions of chains of every length), $\mathcal{M}(\mathsf{ticket}) = \mathbf{Card}_\leq$ (infinite well-ordered cardinals $\aleph_\gamma$ ordered by $\leq$), and $\mathcal{M}(\mathsf{ruler}) = \mathbf{Val}_\leq$ (the thin skeletal poset of infinite valuation carriers $[V]$ generated from discrete cardinals under the covariant valuation operation $[V] \mapsto [\mathbf{2}^V]$, ordered by cardinal injection $|V| \leq |W|$);
- $\mathcal{M}(u_0)(\sigma_\infty) = \omega$, $\mathcal{M}(\pi_{\mathsf{ticket}})(\alpha) = |\alpha|$, $\mathcal{M}(\iota_{\mathsf{init}})(\aleph_\gamma) = \omega_\gamma$, and $\mathcal{M}(\delta_{\mathsf{ruler}})(\aleph_\gamma) = [\aleph_\gamma] \in \mathbf{Val}_\leq$ (with $\mathcal{M}(\delta_{\mathsf{ruler}})(\aleph_0) = \beth_0 := [\mathbb{N}]$ and iterated exponential stages $\beth_{n+1} := [\mathbf{2}^{\beth_n}]$), each preserving $\leq$ functorially and satisfying $\mathcal{M}(\pi_{\mathsf{ticket}}) \circ \mathcal{M}(\iota_{\mathsf{init}}) = \mathrm{id}_{\mathbf{Card}_\leq}$.

The *Grothendieck opfibration of the infinite* is the total category

$$\int_{\mathcal{T}_\infty} \mathcal{M} \;=\; \int_{T \in \mathcal{T}_\infty} \mathcal{M}(T) \xrightarrow{\;p\;} \mathcal{T}_\infty$$

defined by:

- **Objects**: Pairs (T, x) where $T \in \mathrm{Ob}(\mathcal{T}_\infty)$ is a semantic toy and $x \in \mathrm{Ob}(\mathcal{M}(T))$ is a localized configuration in that toy's meaning-fibre.
- **Morphisms**: A morphism $(u, \varphi) : (T, x) \to (T', x')$ consists of a base translation $u : T \to T'$ in $\mathcal{T}_\infty$ together with the unique vertical comparison $\varphi : u_!(x) \leq_{T'} x'$ in $\mathcal{M}(T')$, where $u_! = \mathcal{M}(u)$, composed by $(v, \psi) \circ (u, \varphi) = (v \circ u,\ \psi \circ v_!(\varphi))$.
- **Cocartesian Lifts**: For every object $(T, x) \in \int_{\mathcal{T}_\infty} \mathcal{M}$ and every base morphism $u : T \to T'$ in $\mathcal{T}_\infty$, the pure-translation morphism

$$\widetilde{u}_x \;=\; (u, \mathrm{id}_{u_!(x)}) \;:\; (T, x) \longrightarrow (T', u_!(x))$$

  is p-cocartesian: for any (T'', z) and any morphism $(v \circ u, \psi) : (T, x) \to (T'', z)$ lying over $v \circ u$, there is a unique morphism $(v, \psi) : (T', u_!(x)) \to (T'', z)$ lying over v such that $(v, \psi) \circ \widetilde{u}_x = (v \circ u, \psi)$.
- **Ground Cocartesian Section**: The anchor configurations of the three core toys form a cocartesian section at the countable ground floor:

$$(\mathsf{queue}, \omega) \xrightarrow{\widetilde{\pi}_{\mathsf{ticket}}} (\mathsf{ticket}, \aleph_0) \xrightarrow{\widetilde{\delta}_{\mathsf{ruler}}} (\mathsf{ruler}, \beth_0),$$

  satisfying $\pi_{\mathsf{ticket}!}(\omega) = \aleph_0$, $\iota_{\mathsf{init}!}(\aleph_0) = \omega$, and $\delta_{\mathsf{ruler}!}(\aleph_0) = \beth_0$.

By Thomason's homotopy colimit theorem [28], the classifying nerve of the Grothendieck construction presents the homotopy colimit of the local meaning-fibres:

$$N\Big(\int_{T \in \mathcal{T}_\infty} \mathcal{M}(T)\Big) \;\simeq\; \operatorname*{hocolim}_{T \in \mathcal{T}_\infty} N\big(\mathcal{M}(T)\big).$$




Figure 1. The Grothendieck opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ gluing the semantic toys of the infinite. Left: the Queue Toy fibre $\mathcal{M}(\text{queue}) = \mathbf{Ord}_{\leq}$ climbs via post-limit extensions $(\omega+1, \omega^2, \varepsilon_0)$, all of which map to $\aleph_0$ under the cocartesian pushforward $\pi_{\text{ticket}!}$. Centre: the Ticket Toy fibre $\mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$ climbs via Hartogs initial ordinals $(\aleph_0, \aleph_1, \aleph_2, \ldots, \aleph_\omega)$. Right: the Ruler Toy fibre $\mathcal{M}(\text{ruler}) = \mathbf{Val}_{\leq}$ climbs via the covariant valuation operation $[V] \mapsto [\mathbf{2}^V]$ $(\beth_0, \beth_1, \beth_2)$. The Continuum Hypothesis asks for a model-indexed horizontal bridge $\theta_U : \mathcal{M}_U(\text{ruler}) \dashrightarrow \mathcal{M}_U(\text{ticket})$ mapping $\beth_1$ to a rung of $\mathcal{M}_U(\text{ticket})$: Gödel's L selects $\aleph_1$, Cohen's forcing extensions select $\aleph_2, \aleph_{17}, \ldots$, and König's theorem ($\text{cf}(\beth_1) > \omega$) rules out $\aleph_\omega$.

For our diagram this homotopy type is trivial. The object ruler is terminal in $\mathcal{T}_\infty$: every toy has exactly one composite arrow into it, because $\pi_{\text{ticket}} \circ \iota_{\text{init}} = \text{id}_{\text{ticket}}$. The homotopy colimit is therefore equivalent to $N(\mathcal{M}(\text{ruler}))$, and $\mathcal{M}(\text{ruler})$ has a least element $\beth_0$, so its nerve is contractible:

$$N\Big(\int_{T \in \mathcal{T}_\infty} \mathcal{M}(T)\Big) \simeq *.$$

Passing to the homotopy type inverts every arrow, and in doing so it forgets the directions (escape, absorption, one-way transport) that carry the meaning. This is the thesis of Open Horn Type Theory [20]: meaning-space is not Kan, and its gaps and directions are positive structure that a Kan completion erases. We therefore locate the global meaning of $\sigma_\infty = \infty$ not in the homotopy colimit but in the directed object: the Grothendieck construction $\int_{\mathcal{T}_\infty} \mathcal{M}$ (the lax colimit of $\mathcal{M}$ in **Cat**), together with its projection p, the anchors $x_0(\sigma)$ of the toys and the generative moves $\mathcal{E}_T$ acting on each fibre. The fibres alone would not suffice either: inside a ZFC chart the ruler fibre is order-isomorphic to the ticket fibre through cardinality, and what tells them apart is the move that generates each (Hartogs' move against $V \mapsto \mathbf{2}^V$) and the anchor it starts from. Merging the toys' theories over one shared signature collapses the countable post-limit terms (Theorem 4.1); indexing over $\mathcal{T}_\infty$ keeps them apart, and the directed structure keeps the toys' moves.

Theorem 4.6 (Non-Collapsing Coexistence of Toy Invariants in $\int_{\mathcal{T}_\infty} \mathcal{M}$). *In the Grothendieck opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$:*

(i) The strict inequality $(\text{queue}, \omega) <_{\text{queue}} (\text{queue}, \omega + 1)$ *holds vertically in* $\mathcal{M}(\text{queue}) = \mathbf{Ord}_{\leq}$, *whereas the cocartesian pushforward along* $\pi_{\text{ticket}} : \text{queue} \to \text{ticket}$ *satisfies*

$$\pi_{\text{ticket}!}(\omega) = \aleph_0 = \pi_{\text{ticket}!}(\omega + 1)$$

in $\mathcal{M}(\mathsf{ticket}) = \mathbf{Card}_{\leq}$*, sending the non-identity counit arrow* $\epsilon_{\omega+1} : \omega < \omega + 1$ *in* $\mathbf{Ord}_{\leq}$ *to the identity arrow* $\mathrm{id}_{\aleph_0}$ *in* $\mathbf{Card}_{\leq}$*.*

(ii) Every countable ordinal $\alpha \in [\omega, \omega_1)$ *in the Queue Toy, including* $\omega \cdot 2, \omega^2, \omega^\omega, \varepsilon_0$*, inhabits the vertical inverse-image subcategory* $\pi^{-1}_{\mathsf{ticket!}}(\aleph_0) = [\omega, \omega_1) \subseteq \mathbf{Ord}_{\leq}$*, with* $\iota_{\mathsf{init!}}(\aleph_0) = \omega$ *serving as the strict initial (least) object of* $\pi^{-1}_{\mathsf{ticket!}}(\aleph_0)$*.*

(iii) The dense rational interval $(\mathbb{Q} \cap [0, 1], <)$ *and the natural numbers* $(\mathbb{N}, <)$ *are non-isomorphic in* **LinOrd** *(and* $(\mathbb{Q} \cap [0, 1], <)$ *is not even an object of* $\mathcal{X}_{\mathsf{queue}}$ *since it is ill-founded), while the skip-and-renumber Boustrophedon pairing (2) defines a bijection in* $\mathcal{W}_{\mathsf{ticket}}$ *identifying* $[\mathbb{Q} \cap [0, 1]]_{\mathsf{ticket}} = [\mathbb{Q}]_{\mathsf{ticket}} = \aleph_0$ *in* $\mathcal{M}(\mathsf{ticket})$*.*

Proof. Parts (i) and (ii) follow directly from Proposition 4.4 and Lemma 2.3: in the raw protocol category $\mathcal{X}_{\mathsf{queue}}$, the canonical inclusion $j : \omega \hookrightarrow \omega + 1$ has no reverse order-embedding ($\mathcal{X}_{\mathsf{queue}}(\omega + 1, \omega) = \emptyset$), so $\omega < \omega + 1$ in $\mathbf{Ord}_{\leq}$; under the forgetful functor to $\mathcal{X}_{\mathsf{ticket}}$, Hilbert's shift $b(\omega) = 0, b(n) = n + 1$ provides a reverse injection $\omega + 1 \hookrightarrow \omega$, so $j \in \mathcal{W}_{\mathsf{ticket}}$ and $\pi_{\mathsf{ticket!}}$ maps the counit arrow $\epsilon_\alpha : \omega \leq \alpha$ for every $\alpha \in [\omega, \omega_1)$ to $\mathrm{id}_{\aleph_0}$ in $\mathbf{Card}_{\leq}$ by the triangular identity $(\pi_{\mathsf{ticket!}}\epsilon_\alpha) \circ (\eta_{\pi_{\mathsf{ticket!}}(\alpha)}) = \mathrm{id}_{\aleph_0}$. Because $\pi^{-1}_{\mathsf{ticket!}}(\aleph_0) = [\omega, \omega_1) \subseteq \mathbf{Ord}_{\leq}$ is a thin poset with least element ω, $\epsilon_\alpha : \omega \leq \alpha$ is the unique morphism from ω to α in $\pi^{-1}_{\mathsf{ticket!}}(\aleph_0)$, making ω the strict initial object of the fibre. For Part (iii), the Cantor pairing bijection $J(a, b) = \frac{1}{2}(a + b - 2)(a + b - 1) + a : (\mathbb{N}^+)^2 \xrightarrow{\sim} \mathbb{N}^+$ restricted to reduced coprime pairs $R = \{(a, b) \in (\mathbb{N}^+)^2 \mid \gcd(a, b) = 1\}$ gives an injection $J|_R : \mathbb{Q}_+ \hookrightarrow \mathbb{N}^+$. Composing $J|_R$ with the unique strictly increasing enumeration $\mathrm{rank} : \mathrm{im}(J|_R) \xrightarrow{\sim} \mathbb{N}^+$ of the infinite subset $\mathrm{im}(J|_R) \subseteq \mathbb{N}^+$ (which skips composite-pair indices such as $J(2, 2) = 5$) yields the explicit bijection $\mathrm{ticket} = \mathrm{rank} \circ J|_R : \mathbb{Q}_+ \xrightarrow{\sim} \mathbb{N}^+$ of (2); restricting to $a \leq b$ and adjoining 0 similarly bijects $\mathbb{Q} \cap [0, 1]$ with $\mathbb{N}^+$. □

5 The Three Toy Escalators and Internal Diagonal Engines

We now analyze the three internal escalator endofunctors driving $\mathcal{M}(\mathsf{queue})$, $\mathcal{M}(\mathsf{ticket})$, and $\mathcal{M}(\mathsf{ruler})$, explaining both their categorical mechanics and their intuitive operational meaning.

5.1 The Queue Toy Escalator: Non-Commutative Cocones on $\mathcal{M}(\mathsf{queue})$

In $\mathcal{X}_{\mathsf{queue}} = \mathbf{Ord}_{\mathsf{emb}}$ and its posetal localization $\mathcal{M}(\mathsf{queue}) = \mathbf{Ord}_{\leq}$, why does adding an element on the left ($\mathbf{1} \oplus \omega$) leave the queue unchanged while adding an element on the right ($\omega \oplus \mathbf{1}$) enlarges it?

Categorically, ω equipped with the order-isomorphism $[0, \mathrm{succ}] : \mathbf{1} \oplus \omega \xrightarrow{\sim} \omega$ is the **initial algebra** of the left-successor endofunctor $F(X) = \mathbf{1} \oplus X$ on $\mathbf{Ord}_{\mathsf{emb}}$: for any well-order B with an order-embedding $b : \mathbf{1} \oplus B \hookrightarrow B$, the uniquely defined recursive map $h(0) = b(\star)$, $h(n + 1) = b(h(n))$ is strictly increasing. By **Lambek's Lemma**, the structure map of an initial algebra is an isomorphism, yielding front absorption:

$$\mathbf{1} \oplus \omega \;\cong\; \omega \quad \text{in } \mathcal{X}_{\mathsf{queue}}, \qquad \mathbf{1} \oplus \omega \;=\; \omega \quad \text{in } \mathcal{M}(\mathsf{queue}) = \mathbf{Ord}_{\leq}. \tag{5}$$

Conversely, the right-extension endofunctor $\mathsf{PostLimit}(\alpha) = \alpha \oplus \mathbf{1}$ attaches a **cocone vertex** (a new ceiling $\top$) above the entire infinite chain $0 < 1 < 2 < \dots$ of ω. Every element $n \in \omega$ has only finitely many predecessors $\{0, \dots, n - 1\}$, whereas $\top$ has the entire infinite queue ω ahead of it. The pigeonhole obstruction (Section 3) defeats every candidate order-embedding $\omega \oplus \mathbf{1} \to \omega$:

$$\omega \;<\; \omega \oplus \mathbf{1} \quad \text{in } \mathcal{M}(\mathsf{queue}) = \mathbf{Ord}_{\leq}. \tag{6}$$

Iterating PostLimit and taking suprema (colimits in $\mathbf{Ord}_{\leq}$, corresponding to unions of chains of *initial-segment embeddings* in $\mathbf{Ord}_{\mathsf{emb}}$; note that non-initial shift chains $\omega \xrightarrow{+1} \omega \xrightarrow{+1} \cdots$ do not form initial

segments and have no well-ordered colimit) generates:

$$\omega \xrightarrow{+1} \omega+1 \xrightarrow{+1} \omega+2 \xrightarrow{\text{colim}} \omega\cdot 2 \xrightarrow{\text{colim}} \omega^2 \xrightarrow{\text{colim}} \omega^3 \xrightarrow{\text{colim}} \omega^\omega \xrightarrow{\text{colim}} \varepsilon_0 = \operatorname*{colim}_{n<\omega} \underbrace{\omega^{\omega^{\cdot^{\cdot^{\omega}}}}}_{n}.$$

At Gentzen's ordinal ε_0 [8], the ordinal exponential functor $\alpha \mapsto \omega^\alpha$ reaches its first fixed point ($\omega^{\varepsilon_0} = \varepsilon_0$), while the cocartesian pushforward $\pi_{\text{ticket!}}$ maps the entire chain constantly to $\aleph_0$.

5.2 The Ticket Toy Escalator: Hartogs' Choice-Free Functor on $\mathcal{M}(\text{ticket})$

How does the Ticket Toy $\mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$ climb beyond $\aleph_0$ without borrowing the power-set functor from the Ruler Toy and without assuming the Axiom of Choice?

Look back at Figure 1: sitting vertically above the single ticket size $\aleph_0$ is the entire Queue fibre $\pi^{-1}_{\text{ticket!}}(\aleph_0) = [\omega, \omega_1)$ containing all the distinct ways of lining up $\aleph_0$ elements into a well-ordered queue $(\omega, \omega+1, \omega^2, \varepsilon_0, \dots)$ modulo order-isomorphism. In 1915, Friedrich Hartogs proved in classical ZF (using Power Set, Separation and Replacement, without the Axiom of Choice) that if we collect the ordinal order-type representatives of all well-orderings of subsets of a set X, that collection is itself a well-ordered initial ordinal that cannot inject into X [12]!

Definition 5.1 (Hartogs Ticket Escalator Hartogs). For any well-orderable cardinal $\kappa = |X| \in \mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$ (with initial ordinal representative $X = \omega_\alpha$), the successor ticket size $\text{Hartogs}(\kappa) = \kappa^+$ is the cardinality of the Hartogs initial ordinal:

$$\aleph(X) = \big\{\text{otp}(W, <) \in \text{On} \;\big|\; W \subseteq X \text{ and } < \text{ well-orders } W\big\}.$$

Iterating Hartogs from $\aleph_0 = |\omega|$ and taking suprema at limit ordinals generates the $\aleph$-escalator in $\mathbf{Card}_{\leq}$:

$$\aleph_0,\ \aleph_1 = \text{Hartogs}(\aleph_0),\ \aleph_2 = \text{Hartogs}(\aleph_1),\ \dots,\ \aleph_\omega = \operatorname*{colim}_{n<\omega} \aleph_n,\ \aleph_{\omega+1} = \text{Hartogs}(\aleph_\omega).$$

Proposition 5.2 (Hartogs' Escalator as Fibre Volume of the Queue Opfibration). *In* ZF*, for every well-ordered cardinal $\aleph_\alpha \in \mathcal{M}(\text{ticket}) = \mathbf{Card}_{\leq}$, the initial ordinal $\omega_{\alpha+1} = \iota_{\text{init!}}(\aleph_{\alpha+1})$ of the Hartogs successor $\aleph_{\alpha+1} = \text{Hartogs}(\aleph_\alpha)$ is equal to the cumulative vertical Queue fibre of ordinal order-types of cardinality at most $\aleph_\alpha$:*

$$\iota_{\text{init!}}(\aleph_{\alpha+1}) = \omega_{\alpha+1} = \big\{\beta \in \text{On} \;\big|\; |\beta| \leq \aleph_\alpha\big\},$$

and in the diagnostic game $\mathcal{A}_{\text{ticket}}(\aleph_\alpha, \aleph_{\alpha+1})$, the Challenger possesses an explicit choice-free winning obstruction against every Proposer injection $f : \omega_{\alpha+1} \hookrightarrow \omega_\alpha$, establishing $\aleph_\alpha <_{\text{ticket}} \aleph_{\alpha+1}$.

Proof. By Definition 5.1, an ordinal $\beta \in \text{On}$ belongs to $\aleph(\omega_\alpha)$ if and only if $\beta = \text{otp}(W, <)$ for some well-ordered subset $W \subseteq \omega_\alpha$, which holds if and only if $|\beta| \leq |\omega_\alpha| = \aleph_\alpha$. Since ω_α is an ordinal, the inclusion $\omega_\alpha \hookrightarrow \omega_{\alpha+1}$ witnesses $\aleph_\alpha \preceq_{\text{ticket}} \aleph_{\alpha+1}$. Conversely, if a Proposer presents an injective map $f : \omega_{\alpha+1} \hookrightarrow \omega_\alpha$, then transporting the canonical well-ordering of $\omega_{\alpha+1}$ onto its image $W = \text{im}(f) \subseteq \omega_\alpha$ exhibits $\omega_{\alpha+1}$ as the order type $\text{otp}(W, f_*(\in))$ of a well-ordered subset of ω_α. By definition of $\aleph(\omega_\alpha) = \omega_{\alpha+1}$, this forces $\omega_{\alpha+1} \in \omega_{\alpha+1}$ (i.e., $\omega_{\alpha+1} < \omega_{\alpha+1}$), an irreflexive contradiction. Thus the Challenger refutes every candidate injection f via $\chi(f) = \text{otp}(\text{im}(f), f_*(\in))$, proving $\aleph_\alpha <_{\text{ticket}} \aleph_{\alpha+1}$ in ZF. □

5.3 The Ruler Toy Escalator: Lawvere's Theorem as Uniform Challenger Obstruction

In the Ruler Toy fibre $\mathcal{M}(\text{ruler}) = \mathbf{Val}_{\leq}$, escalation is governed by the covariant valuation operation $[V] \mapsto [\mathbf{2}^V]$. Before stating Lawvere's theorem, let us clarify three categorical concepts for philosophical readers:

(a) **What is the Truth-Value Object Ω?** In set theory, we speak of "subsets" $S \subseteq A$. In categorical logic, a subset or binary colouring on a domain A is represented by its characteristic valuation map $c : A \to \Omega$ into a truth-value object Ω. In classical two-valued logic, $\Omega = \mathbf{2} = \{0, 1\} \cong \{L, R\}$, where each element $a \in A$ is assigned one of two values.

(b) **What is a Cartesian Closed Category (CCC) and Currying?** Ambiently in **Set** or any topos $\mathcal{E}$, the category is *cartesian closed*: for any two objects A and Ω, the collection of all valuation morphisms $A \to \Omega$ forms an exponential object Ω^A. **Currying** translates bijectively between a catalogue of A-many valuations $e : A \to \Omega^A$ and a two-dimensional evaluation grid $\widehat{e} : A \times A \to \Omega$, where $\widehat{e}(a_1, a_2) = e(a_1)(a_2)$ is the value that the a_1-th valuation assigns to locus a_2. (While $\Omega^{(-)}$ is contravariant on arbitrary maps $f : A \to B$ via precomposition $q \mapsto q \circ f$, on injective carrier embeddings $i : A \hookrightarrow B$ in $\mathcal{X}_{\mathsf{ruler}}$ it acts *covariantly* by zero-extension $i_! : \mathbf{2}^A \hookrightarrow \mathbf{2}^B$, inducing the monotone endofunctor $[V] \mapsto [\mathbf{2}^V]$ on $\mathcal{M}(\mathsf{ruler}) = \mathbf{Val}_{\leq}$.)

(c) **What is Lawvere's Diagonal Operator?** In 1969, F. William Lawvere showed that Cantor's 1891 diagonal argument is a purely structural consequence of currying [16]: given any Proposer catalogue $e : A \to \Omega^A$, the Challenger walks down the main diagonal of the 2D table via $\Delta_A : a \mapsto (a, a)$, reads the value $\widehat{e}(a, a)$ that the a-th valuation assigns to a itself, and applies a fixed-point-free flip $\sigma : \Omega \to \Omega$ ($0 \leftrightarrow 1$).

Within our framework (Definition 2.1), Lawvere's construction defines the explicit **uniform winning Challenger obstruction operator** χ_{Lawvere} in the diagnostic game $\mathcal{A}_{\mathsf{ruler}}(A, \Omega^A)$.

Theorem 5.3 (Lawvere's Diagonal Obstruction and Strict Valuation Escape). *(i)* ***General Categorical Diagonal Obstruction****: Let $\mathcal{E}$ be a cartesian closed category, let $A, \Omega \in \mathrm{Ob}(\mathcal{E})$, and let $\sigma : \Omega \to \Omega$ be an endomorphism with no fixed point on global elements (no global element $y : \mathbf{1} \to \Omega$ satisfies $\sigma \circ y = y$). Define the Challenger obstruction operator*

$$\chi_{\text{Lawvere}} : \mathcal{E}(A, \Omega^A) \longrightarrow \mathcal{E}(\mathbf{1}, \Omega^A)$$


by assigning to each Proposer catalogue $e : A \to \Omega^A$ the curried transpose $\ulcorner d_e \urcorner : \mathbf{1} \to \Omega^A$ of the diagonal morphism

$$d_e \;=\; \sigma \circ \widehat{e} \circ \Delta_A \;:\; A \xrightarrow{\Delta_A} A \times A \xrightarrow{\widehat{e}} \Omega \xrightarrow{\sigma} \Omega \tag{7}$$

(where $\Delta_A = \langle \mathrm{id}_A, \mathrm{id}_A \rangle$ and $\widehat{e} : A \times A \to \Omega$ is the uncurried evaluation of e). Then for every Proposer morphism $e : A \to \Omega^A$ and every global point $a_0 : \mathbf{1} \to A$, we have $e \circ a_0 \neq \chi_{\text{Lawvere}}(e)$ (no $e : A \to \Omega^A$ is weakly point-surjective).

(ii) ***Strict Generative Escape ($[A] <_{\mathsf{ruler}} [\mathbf{2}^A]$)****: For any non-empty carrier A in* **Set** *with $\Omega = \mathbf{2} = \{0, 1\}$ and negation $\sigma(x) = 1 - x$, the singleton-indicator map $\{\cdot\}_A : A \hookrightarrow \mathbf{2}^A$ ($a \mapsto \delta_a$) provides a forward embedding $A \preceq_{\mathsf{ruler}} \mathbf{2}^A$, while Part (i) ensures that no surjection $e : A \twoheadrightarrow \mathbf{2}^A$ (and hence no reverse injection $\mathbf{2}^A \hookrightarrow A$) exists, establishing strict generative escape $[A] <_{\mathsf{ruler}} [\mathbf{2}^A]$ in $\mathcal{M}(\mathsf{ruler}) = \mathbf{Val}_{\leq}$.*

Proof. (i) Suppose toward a contradiction that for some $e : A \to \Omega^A$ there exists a global element $a_0 : \mathbf{1} \to A$ such that $e \circ a_0 = \ulcorner d_e \urcorner$, or equivalently $\widehat{e} \circ (a_0 \times \mathrm{id}_A) = d_e$. Evaluating both sides at a_0 via the diagram



gives

$$\widehat{e}(a_0, a_0) \;=\; d_e(a_0) \;=\; \sigma\big(\widehat{e}(\Delta_A(a_0))\big) \;=\; \sigma\big(\widehat{e}(a_0, a_0)\big),$$

exhibiting $y = \widehat{e}(a_0, a_0) : \mathbf{1} \to \Omega$ as a global fixed point of σ, contradicting the hypothesis on σ. (ii) Since $0 \neq 1$ in $\mathbf{2}$, $\sigma(x) = 1 - x$ has no fixed point, and $\{\cdot\}_A(a)(b) = 1 \iff a = b$ injects A into $\mathbf{2}^A$.

In **Set** a surjection $A \twoheadrightarrow \mathbf{2}^A$ is point-surjective, which Part (i) excludes. (In other toposes, Part (i) still excludes point-surjections; passing from there to a strict cardinal comparison needs further hypotheses, which we do not use.) If an injection $g : \mathbf{2}^A \hookrightarrow A$ existed, its retraction $e : A \twoheadrightarrow \mathbf{2}^A$ ($e \circ g = \mathrm{id}_{\mathbf{2}^A}$, defined as g^{-1} on the image of g and constant on its complement) would yield a point-surjective map $e : A \to \mathbf{2}^A$, contradicting Part (i). Thus $[A] <_{\mathsf{ruler}} [\mathbf{2}^A]$ in $\mathbf{Val}_{\leq}$. □

At Stage 1 ($\beth_1$), $A = \mathbb{N}$, $\Omega = \{L, R\}$, and $\sigma =$ flip; at Stage 2 ($\beth_2$), $A = [0,1]$, $\Omega = \mathbf{2}$, and $\sigma(c) = 1 - c$. Iterating $V_{\alpha+1} = \Omega^{V_\alpha}$ generates the strict hierarchy $\beth_0 <_{\mathsf{ruler}} \beth_1 <_{\mathsf{ruler}} \beth_2 <_{\mathsf{ruler}} \cdots$ inside $\mathcal{M}(\mathsf{ruler})$ independently of Hartogs.

6 The Continuum Hypothesis as Model-Indexed Horizontal Bridge and Base-Change Cocycle

6.1 Model-Indexed Horizontal Bridges and the Pair-Groupoid Shift Cocycle

In the Grothendieck opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$, the two terms of the Continuum Hypothesis inhabit distinct rooms:

(i) $(\mathsf{ticket}, \aleph_1)$ is the first successor object in $\mathcal{M}(\mathsf{ticket}) = \mathbf{Card}_{\leq}$ under the extensional escalator Hartogs;

(ii) $(\mathsf{ruler}, \beth_1)$ is the first exponential object $[\mathbf{2}^{\beth_0}]$ in $\mathcal{M}(\mathsf{ruler}) = \mathbf{Val}_{\leq}$ under the intensional valuation escalator;

(iii) $p(\mathsf{ticket}, \aleph_1) = \mathsf{ticket} \neq \mathsf{ruler} = p(\mathsf{ruler}, \beth_1)$.

Let us pause to appreciate what this means both mathematically and philosophically. Inside the Ticket Toy T_{ticket}, we construct $\aleph_1$ by collecting countable queues, and inside the Ruler Toy T_{ruler}, we construct $\beth_1$ by taking binary cuts over $\beth_0$. Both toys start from the same ground floor ($\delta_{\mathsf{ruler}!}(\aleph_0) = \beth_0$). However, **the base category $\mathcal{T}_\infty$ contains no morphism** ruler $\to$ ticket. (Note a point of categorical precision: in fibred category theory [10, 2], an internal *cleavage* of $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ chooses cocartesian lifts over arrows *already present* in $\mathcal{T}_\infty$; because $\mathrm{Hom}_{\mathcal{T}_\infty}(\mathsf{ruler}, \mathsf{ticket}) = \emptyset$, comparing $\beth_1$ with the $\aleph$-escalator is not an internal lift over $\mathcal{T}_\infty$, but requires *extending the base* inside an ambient set-theoretic or topos model U.)

Specifically, a **bridging meta-toy** U (a countable transitive model of ZFC) supplies a **model-indexed horizontal bridge functor (cardinal-evaluation functor)**

$$\theta_U : \mathcal{M}_U(\mathsf{ruler}) \longrightarrow \mathcal{M}_U(\mathsf{ticket}), \qquad [V]_{\mathsf{ruler}} \longmapsto |V|^U,$$

which invokes the Axiom of Choice in U to well-order each valuation carrier $V \in \mathcal{M}_U(\mathsf{ruler})$ and read off its initial cardinal rung $\theta_U(\beth_1^U) = \aleph_{\gamma(U)}^U$ in $\mathcal{M}_U(\mathsf{ticket})$. While $\gamma(U)$ is uniquely determined inside any single fixed ZFC model U, it varies when we pass between models $U \rightsquigarrow U'$ via forcing extensions.

Definition 6.1 (Continuum Bridge Index and Pair-Groupoid Shift Cocycle). Let $\mathcal{S}$ be the category of bridging meta-toy charts, whose objects $U \in \mathcal{S}$ are countable transitive models of ZFC and whose morphisms $f : U \rightsquigarrow U'$ are countable-chain-condition (**ccc**) forcing extensions $U' = U[G]$ (where the generic filter G is a point of the Boolean sheaf topos $\mathbf{Sh}^U(\mathcal{B}(\mathbb{P}))$, through which $U[G]$ interprets the Boolean-valued names of U; the global-sections morphism $\mathbf{Sh}^U(\mathcal{B}(\mathbb{P})) \to \mathbf{Set}^U$ is fixed by $\mathbb{P}$ alone and does not depend on G). We work under the hypothesis that a countable transitive model of ZFC exists, which is strictly stronger than Con(ZFC); the relative-consistency results themselves can be obtained from Con(ZFC) alone by Boolean-valued or syntactic forcing, which we do not reproduce. For each chart $U \in \mathcal{S}$:

(i) The *continuum bridge index* is the ordinal $\gamma(U) \geq 1$ in On^U defined by

$$\theta_U(\beth_1^U) \;=\; \left|\mathbf{2}^{\aleph_0}\right|^U \;=\; \aleph_{\gamma(U)}^U \qquad \text{in } \mathcal{M}_U(\mathsf{ticket}), \tag{8}$$

and more generally, for each ordinal β, the *beth bridge index* $\gamma_{\beth}(U,\beta)$ is defined by $\theta_U(\beth_\beta^U) = \aleph^U_{\gamma_{\beth}(U,\beta)}$. The *Continuum Hypothesis (CH)* holds in U if and only if $\gamma(U) = 1$, and the *Generalized Continuum Hypothesis (GCH)* implies $\gamma_{\beth}(U,\beta) = \beta$ for all ordinals β.

(ii) Because every ccc forcing extension $f : U \rightsquigarrow U'$ preserves ordinals and cardinals ($\mathrm{On}^{U'} = \mathrm{On}^U$ and $\aleph_\alpha^{U'} = \aleph_\alpha^U$), the variation of the horizontal bridge $\theta_U(\beth_1)$ across f is recorded by the *shift 1-cocycle* with values in the pair groupoid $\mathrm{Pair}(\mathrm{On})$:

$$\Delta\gamma(f) \;:=\; \big(\gamma(U),\,\gamma(U')\big) \;\in\; \mathrm{Hom}_{\mathrm{Pair}(\mathrm{On})}\big(\gamma(U),\,\gamma(U')\big), \tag{9}$$

which satisfies the functorial cocycle identity $\Delta\gamma(g \circ f) = \Delta\gamma(g) \circ \Delta\gamma(f)$ for composable ccc extensions $U \xrightarrow{f} U' \xrightarrow{g} U''$, and is non-trivial ($\Delta\gamma(f) \neq \mathrm{id}$) whenever $\gamma(U') \neq \gamma(U)$.

In a pair groupoid every 1-cocycle is a coboundary, so $\Delta\gamma$ carries no invariant beyond the pair of indices: it records that the bridge moved, and from where to where. We use it as bookkeeping for comparisons across models, not as a cohomological invariant.

6.2 Why the Bridge Slides: Global Sections of Ω vs. Rigidity of $\aleph$

Why does the horizontal bridge $\theta_U(\beth_1)$ slide across the $\aleph$-ladder when we pass from V to a forcing extension $V[G]$? Sheaf and topos theory (Lawvere, Tierney, and Mac Lane–Moerdijk [18]) illuminates the exact structural mechanism when we distinguish the internal logic of the Boolean sheaf topos $\mathbf{Sh}(\mathcal{B}(\mathbb{P}))$ from its external global sections:

1. **Internal $\Omega \cong \mathbf{1}+\mathbf{1}$ vs. External Global Truth-Values $\Gamma(\Omega) \cong \mathcal{B}(\mathbb{P})$:** Given a forcing poset $\mathbb{P} \in V$, let $\mathcal{B} = \mathcal{B}(\mathbb{P}) \in V$ be its regular-open complete Boolean algebra, and let $f : \mathbf{Sh}(\mathcal{B}) \to \mathbf{Set}^V$ be the canonical geometric morphism ($f^* \dashv f_*$). Because $\mathbf{Sh}(\mathcal{B})$ is a Boolean topos, its *internal* subobject classifier is isomorphic to the binary coproduct $\Omega_{\mathbf{Sh}(\mathcal{B})} \cong \mathbf{1}+\mathbf{1} \cong f^*(\mathbf{2}_V)$. However, when $\mathcal{B}$ is non-trivial, $\mathbf{Sh}(\mathcal{B})$ is not well-pointed: the terminal object decomposes in many ways, so $\Gamma(\mathbf{1}+\mathbf{1})$ is not the two-element set, and the **external algebra of global truth-values (global sections of Ω)** expands from $\{0,1\}$ to the entire complete Boolean algebra $\mathcal{B}$:

$$\Gamma\big(\Omega_{\mathbf{Sh}(\mathcal{B})}\big) \;:=\; \mathrm{Hom}_{\mathbf{Sh}(\mathcal{B})}(\mathbf{1},\Omega_{\mathbf{Sh}(\mathcal{B})}) \;\cong\; \mathrm{Sub}_{\mathbf{Sh}(\mathcal{B})}(\mathbf{1}) \;\cong\; \mathcal{B}(\mathbb{P}).$$

2. **Non-Preservation of the Exponential $\Omega^{f^*(\omega)}$:** While the inverse-image functor f^* preserves the finite coproduct $\mathbf{2} = \mathbf{1}+\mathbf{1}$, f^* **does not preserve infinite exponential objects** when $\mathbb{P}$ adds new reals:

$$f^*\big(\mathbf{2}_V^{\omega}\big) \;\subsetneq\; \Omega_{\mathbf{Sh}(\mathcal{B})}^{f^*(\omega)}.$$

Externally, a global section of $\Omega_{\mathbf{Sh}(\mathcal{B})}^{f^*(\omega)}$ is a morphism $f^*(\omega) \to \Omega_{\mathbf{Sh}(\mathcal{B})}$, which corresponds to a Boolean-valued name $\tau : \omega \to \mathcal{B}(\mathbb{P})$ assigning an independent Boolean truth-value $[\![n \in \tau]\!] \in \mathcal{B}(\mathbb{P})$ to each $n \in \omega$. Interpreting these names through a V-generic filter G evaluates them to genuinely new subsets of ω in $V[G]$.

3. **Why the Ticket Toy's $\aleph$-Ladder Stays Rigid Under the ccc:** When $\mathbb{P}$ satisfies the **countable chain condition (ccc)** in V—meaning every antichain of mutually incompatible conditions in $\mathbb{P}$ is at most countable in V—a name for the value of a function at any given ordinal has at most countably many possible values in V. Consequently, no uncountable well-ordered initial ordinal can be collapsed or have its cofinality reduced: **the Ticket Toy's cardinal spine $\mathrm{Card}_{\leq}$ is invariant between V and $V[G]$** ($\aleph_\alpha^{V[G]} = \aleph_\alpha^V$), whereas the Ruler Toy's continuum $\beth_1^{V[G]} = (\mathbf{2}^{\omega})^{V[G]}$ absorbs the generic Boolean cuts!

Theorem 6.2 (Differential Variance of the Ticket and Ruler Toys under Forcing). *Let V be a transitive model of* ZFC + GCH *(for example a countable transitive model of $V = L$, in which $\gamma = 1$), and let $\mathbb{P} \in V$ be a forcing poset satisfying the countable chain condition (**ccc**) in V, with generic extension $V[G]$. Then:*

(i) ***Rigidity of the Ticket Toy Fibre*** $\mathcal{M}(\mathsf{ticket}) = \mathbf{Card}_{\leq}$*: Every infinite cardinal and its cofinality are preserved between* V *and* $V[G]$ *(*$\mathrm{cf}^{V[G]}(\lambda) = \mathrm{cf}^{V}(\lambda)$ *and* $\aleph_\alpha^{V[G]} = \aleph_\alpha^V$ *for all ordinals* α*).*

(ii) ***Elasticity of the Ruler Toy Fibre*** $\mathcal{M}(\mathsf{ruler}) = \mathbf{Val}_{\leq}$ ***Under Cohen Forcing****: For the Cohen poset* $\mathbb{P}_\beta = \mathrm{Fn}(\aleph_\beta \times \omega, 2) \in V$ *with* $\mathrm{cf}^V(\aleph_\beta) > \omega$*, the generic extension* $V[G]$ *adjoins* $\aleph_\beta$ *generic binary cuts, yielding*

$$\theta_{V[G]}\big(\beth_1^{V[G]}\big) \;=\; \aleph_\beta, \qquad \Delta\gamma(f_\beta) \;=\; (1,\,\beta),$$

which preserves CH *when* $\beta = 1$ *(*$\Delta\gamma(f_1) = \mathrm{id}_1$*) and produces a strict cross-fibre shift* $\Delta\gamma(f_\beta) \neq \mathrm{id}$ *for every* $\beta \geq 2$ *with* $\mathrm{cf}^V(\aleph_\beta) > \omega$*.*

Proof. (i) To prove cardinal and cofinality preservation, suppose a condition $p \in \mathbb{P}$ forces that $\dot{g} : \check{\delta} \to \check{\lambda}$ is a function in $V[G]$ from an infinite ordinal δ into λ. For each $\alpha < \delta$, define in V the set of possible values

$$A_\alpha \;=\; \big\{\xi < \lambda \;\big|\; \exists q \leq p,\; q \Vdash \dot{g}(\check{\alpha}) = \check{\xi}\big\}.$$

Using the Axiom of Choice in V, choose for each $\xi \in A_\alpha$ a witness $q_{\alpha,\xi} \leq p$ forcing $\dot{g}(\check{\alpha}) = \check{\xi}$. Since $\dot{g}$ is forced to be a function, $\{q_{\alpha,\xi} \mid \xi \in A_\alpha\}$ is an antichain in $\mathbb{P}$, which is countable in V by the ccc ($|A_\alpha|^V \leq \aleph_0$). Thus $\dot{g}$ is forced to have its range contained in the ground-model set $A = \bigcup_{\alpha<\delta} A_\alpha \in V$ of cardinality $|A|^V \leq |\delta|^V \cdot \aleph_0 = |\delta|^V$. If $\delta = \kappa < \lambda$ for cardinals κ, λ in V, then $|A|^V \leq \kappa < \lambda$, so $\dot{g}$ cannot be surjective onto λ; hence cardinals are preserved. Likewise, if λ is a limit ordinal of uncountable cofinality $\mathrm{cf}^V(\lambda) > \omega$ and $\delta < \mathrm{cf}^V(\lambda)$, then $|A|^V \leq |\delta|^V \cdot \aleph_0 < \mathrm{cf}^V(\lambda)$, so A is bounded below λ and $\dot{g}$ cannot be cofinal in λ (while if $\mathrm{cf}^V(\lambda) = \omega$, $\mathrm{cf}^{V[G]}(\lambda) = \omega$ since no limit ordinal has finite cofinality). Thus all cardinals and cofinalities are preserved.

(ii) Given a V-generic filter $G \subseteq \mathrm{Fn}(\aleph_\beta \times \omega, 2)$, the union $\bigcup G : \aleph_\beta \times \omega \to 2$ curries into $\aleph_\beta$ binary sequences $c_\xi : \omega \to 2$ ($\xi < \aleph_\beta$). For $\xi \neq \eta$, the set $D_{\xi,\eta} = \{p \in \mathbb{P}_\beta \mid \exists n < \omega,\, p(\xi, n) \neq p(\eta, n)\}$ is dense in $\mathbb{P}_\beta$, so $c_\xi \neq c_\eta$ in $V[G]$ and $\aleph_\beta \leq |\beth_1|^{V[G]}$. Conversely, every subset of ω in $V[G]$ has a nice $\mathbb{P}_\beta$-name determined by ω-many antichains of $\mathbb{P}_\beta$. Since $|\mathbb{P}_\beta|^V = \aleph_\beta$ and every antichain is countable by the ccc, the number of nice names in V is at most $\big((\aleph_\beta^{\aleph_0})^{\aleph_0}\big)^V = (\aleph_\beta^{\aleph_0})^V$. In $V \models \mathrm{GCH}$, $\mathrm{cf}^V(\aleph_\beta) > \omega$ implies $(\aleph_\beta^{\aleph_0})^V = \aleph_\beta$, whence $|\beth_1|^{V[G]} = \aleph_\beta$ and $\gamma(V[G]) = \beta$. □

6.3 König's Obstruction at $\aleph_\omega$ as a Cross-Fibre Conflict

Although the bridge index $\gamma(U)$ varies across $\mathcal{S}$ via $\Delta\gamma(f)$, König's theorem [13] forbids $\gamma(U) = \omega$ ($\theta_U(\beth_1) \neq \aleph_\omega$). In the Grothendieck opfibration $\int_{\mathcal{T}_\infty} \mathcal{M}$, this obstruction has a transparent meaning:

- In the Ticket Toy $\mathcal{M}(\mathsf{ticket})$, $\aleph_\omega = \mathrm{colim}_{n<\omega} \aleph_n$ is the supremum of a countable sequence of smaller rungs $\aleph_0 < \aleph_1 < \aleph_2 < \dots$, which allows the Challenger to run a block-diagonal argument across the ω-many rungs and prove $\aleph_\omega < \aleph_\omega^{\aleph_0}$.
- In the Ruler Toy $\mathcal{M}(\mathsf{ruler})$, however, $\beth_1 = [\mathbf{2}^{\mathbb{N}}]$ is already an infinite product of copies of $\mathbf{2}$ indexed by $\mathbb{N}$. Because a grid of $\mathbb{N}\times\mathbb{N}$ tickets is bijective with a single row of $\mathbb{N}$ tickets via the Boustrophedon snake-path (2), taking a countable product of copies of the Ruler $\beth_1$ leaves its cardinality unchanged ($\beth_1^{\aleph_0} = \beth_1$).

Therefore, $\beth_1$ can never bridge onto $\aleph_\omega$: one is invariant under countable products while the other is strictly enlarged by them!

Theorem 6.3 (König's Cross-Fibre Cofinality Obstruction). *In every* ZFC *meta-toy chart* $U \in \mathcal{S}$*, the bridge image* $\theta_U(\beth_1^U) \in \mathcal{M}_U(\mathsf{ticket})$ *satisfies*

$$\mathrm{cf}\big(\theta_U(\beth_1^U)\big) \;>\; \omega. \tag{10}$$

In particular, for any cardinal $\kappa = \mathrm{colim}_{n<\omega} \kappa_n$ *of countable cofinality in* $\mathcal{M}_U(\mathsf{ticket})$ *(such as* $\aleph_\omega = \mathrm{colim}_{n<\omega} \aleph_n$*), no chart* $U \in \mathcal{S}$ *satisfies* $\theta_U(\beth_1^U) = \kappa$*.*

Proof. Work inside any ZFC chart $U \in \mathcal{S}$.

Step 1. **Countable product invariance of $\mathbf{2}^{\mathbb{N}}$**: Since $\mathbb{N} \times \mathbb{N} \cong \mathbb{N}$ via the bijection (2), currying in $\mathbf{Set}^U$ gives the bijection

$$\left(\mathbf{2}^{\mathbb{N}}\right)^{\mathbb{N}} \;\cong\; \mathbf{2}^{\mathbb{N}\times\mathbb{N}} \;\cong\; \mathbf{2}^{\mathbb{N}},$$

which maps under $\theta_U([V]) = |V|^U$ to the cardinal equation

$$\theta_U(\beth_1)^{\aleph_0} \;=\; \theta_U(\beth_1). \tag{11}$$

Step 2. **Diagonal Challenger obstruction over countable colimits in $\mathcal{M}_U(\text{ticket})$**: Suppose $\kappa = \sup_{n<\omega} \kappa_n$ with $\kappa_0 < \kappa_1 < \cdots < \kappa$. Given any Proposer map $f : \kappa \to \prod_{n<\omega} \kappa_{n+1}$, partition $\kappa = \bigsqcup_{n<\omega} A_n$ with $|A_n| = \kappa_n$. Since $|\pi_n(f(A_n))| \le \kappa_n < \kappa_{n+1}$, the Challenger selects the least ordinal $d_n \in \kappa_{n+1} \setminus \pi_n(f(A_n))$ for each $n < \omega$. The diagonal tuple $d = (d_0, d_1, \dots) \in \prod_{n<\omega} \kappa_{n+1} \subseteq \kappa^{\aleph_0}$ differs from $f(\alpha)$ at coordinate n for all $\alpha \in A_n$, proving

$$\kappa \;=\; \sum_{n<\omega} \kappa_n \;<\; \prod_{n<\omega} \kappa_{n+1} \;\le\; \kappa^{\aleph_0}. \tag{12}$$

Step 3. **Conclusion**: Combining (11) and (12), any identification $\theta_U(\beth_1) = \kappa$ with $\mathrm{cf}(\kappa) = \omega$ would yield $\kappa < \kappa^{\aleph_0} = \kappa$, a contradiction. Thus $\mathrm{cf}(\theta_U(\beth_1)) > \omega$. □

By Easton's theorem [6], monotonicity ($\kappa \le \lambda \implies 2^\kappa \le 2^\lambda$) and König's cofinality bound ($\mathrm{cf}(2^\kappa) > \kappa$) are the only ZFC-provable constraints on the continuum function restricted to regular cardinals: if $V \models \mathrm{GCH}$ and F is a class function of V, definable there, from the regular cardinals to cardinals, monotone and with $\mathrm{cf}(F(\kappa)) > \kappa$, then a cofinality-preserving class forcing yields an extension satisfying ZFC in which $2^\kappa = F(\kappa)$ for every regular κ. At singular cardinals ZFC proves further constraints, which we do not use.

7 A Type-Theoretic Sketch and a Worked Comparative Calculus

The type-theoretic part of this section is a conceptual sketch, not a formalization: we have not checked it in a proof assistant, and the steps that need the law of excluded middle or propositional resizing are marked. The singleton map $A \to (A \to \mathbf{2})$ and zero-extension along an injection are classical too: constructively they need decidable equality on A and decidable membership in the image.

7.1 Why Directed Precategories Are Needed in Homotopy Type Theory

In modern constructive and univalent foundations such as Homotopy Type Theory (HoTT) [30] and its temporal and open-horn extensions for variable sense (DHoTT [19] and OHTT [20]; see also [24, 25, 26]), it is natural to ask how the Grothendieck opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$ is represented as a formal type.

Here an instructive subtlety arises that illuminates the directed nature of our semantic toys:

- In standard HoTT, the built-in identity type $a =_A b$ is automatically **symmetric** (every path $p : a =_A b$ has an inverse $p^{-1} : b =_A a$). If we tried to represent translations between toys as identity paths $p : \mathsf{queue} =_{\mathsf{Toy}} \mathsf{ticket}$, then every translation between toys would be invertible, which would collapse the Queue Toy and the Ticket Toy into the same fibre! Worse still, if we defined $\mathsf{Toy}_0 = \{\mathsf{queue}, \mathsf{ticket}, \mathsf{ruler}\}$ as a discrete 3-element enumeration type, then the identity type $\mathsf{queue} =_{\mathsf{Toy}_0} \mathsf{ticket}$ would be empty (0), forbidding any transport between toys at all.
- Our inter-toy corridors ($\pi_{\mathsf{ticket}} : \mathsf{queue} \to \mathsf{ticket}$ and $\delta_{\mathsf{ruler}} : \mathsf{ticket} \to \mathsf{ruler}$) are **directed, one-way functors**: you can forget the waiting-line order on $\omega + 1$ to obtain $\aleph_0$ tickets, but you cannot uniquely recover the waiting line $\omega + 1$ from the bare ticket count $\aleph_0$.

Therefore, following Ahrens, Kapulkin, and Shulman [1] and the fibred type-theoretic semantics of [19, 20, 24], we represent the base $\mathcal{T}_\infty$ inside HoTT as a **1-truncated precategory** (a type of objects $\mathrm{Ob}_{\mathcal{T}_\infty}$ equipped with an explicit family of directed hom-sets $\mathrm{Hom}_{\mathcal{T}_\infty}(T_1, T_2)$), and we represent the thin

meaning-fibres $\mathcal{M}(T)$ as **posetal precategories** (whose hom-types are (-1)-truncated propositions, [1, Example 3.14]) over a common univalent universe $\mathcal{U}_0$ so that all transport functors are defined on the full object ranges.

Definition 7.1 (Type-Theoretic Specification of the Opfibration $\int_{\mathcal{T}_\infty} \mathcal{M}$). Fix nested univalent universes $\mathcal{U}_0 : \mathcal{U}_1$.

(i) **The Base Precategory $\mathcal{T}_\infty$**: Let $\mathsf{Ob}_{\mathcal{T}_\infty} \equiv \{\mathsf{queue}, \mathsf{ticket}, \mathsf{ruler}\} : \mathcal{U}_0$, and define the directed hom-sets $\mathsf{Hom}_{\mathcal{T}_\infty} : \mathsf{Ob}_{\mathcal{T}_\infty} \to \mathsf{Ob}_{\mathcal{T}_\infty} \to \mathsf{hSet}_{\mathcal{U}_0}$ generated by $\pi_{\mathsf{ticket}} : \mathsf{Hom}_{\mathcal{T}_\infty}(\mathsf{queue}, \mathsf{ticket})$, $\iota_{\mathsf{init}} : \mathsf{Hom}_{\mathcal{T}_\infty}(\mathsf{ticket}, \mathsf{queue})$, and $\delta_{\mathsf{ruler}} : \mathsf{Hom}_{\mathcal{T}_\infty}(\mathsf{ticket}, \mathsf{ruler})$, subject to the relation $\pi_{\mathsf{ticket}} \circ \iota_{\mathsf{init}} = \mathsf{id}_{\mathsf{ticket}}$.

(ii) **The Thin Fibered Precategories $\mathcal{M}(T) = (\mathcal{M}_0(T), \mathsf{Hom}_{\mathcal{M}(T)})$ for $T : \mathsf{Ob}_{\mathcal{T}_\infty}$:**

- $\mathcal{M}_0(\mathsf{queue}) \equiv \left\|\mathsf{Ord}_{\mathcal{U}_0}\right\|_0$, the 0-truncation of the univalent type $\mathsf{Ord}_{\mathcal{U}_0}$ of well-ordered sets $(A, <_A)$ in $\mathcal{U}_0$ [30, §10.3], equipped with the (-1)-truncated hom-proposition $\mathsf{Hom}_{\mathcal{M}(\mathsf{queue})}(\|(A, <_A)\|_0, \|(B, <_B)\|_0) \equiv \|(A, <_A) \hookrightarrow_{\mathsf{ord}} (B, <_B)\|_{-1}$ of mere order-preserving embedding (the inductive Brouwer–Cantor tree type Brw with constructors zero, succ, $\mathsf{lim}_{\mathbb{N}}$ maps into $\mathcal{M}_0(\mathsf{queue})$ by sending a tree to the ordinal it denotes; the map is not injective, since distinct trees can denote the same ordinal, so the constructive countable sub-fibre is an extensional quotient of Brw, as compared in [14]).
- $\mathcal{M}_0(\mathsf{ticket}) \equiv \left\|\mathsf{Card}_{\mathcal{U}_0}\right\|_0$, the 0-truncation of the type $\mathsf{Card}_{\mathcal{U}_0} \equiv \Sigma(X : \mathsf{hSet}_{\mathcal{U}_0}).\, \|\mathsf{isWellOrderable}(X)\|_{-1}$ of well-orderable sets in $\mathcal{U}_0$ (whose identity type is mere bijection $\|X \simeq Y\|_{-1}$, by Univalence and [30, Thm. 7.3.12], so $\mathbb{N} \sqcup \mathbf{1} \simeq \mathbb{N}$ yields $\aleph_0 + 1 =_{\mathcal{M}_0(\mathsf{ticket})} \aleph_0$), equipped with the hom-proposition $\mathsf{Hom}_{\mathcal{M}(\mathsf{ticket})}(\|X\|_0, \|Y\|_0) \equiv \|X \hookrightarrow Y\|_{-1}$ of mere injection.
- $\mathcal{M}_0(\mathsf{ruler}) \equiv \left\|\mathsf{Val}_{\mathcal{U}_0}\right\|_0$, the 0-truncation of the carrier family $\mathsf{Val}_{\mathcal{U}_0} : \mathcal{U}_1$ inductively generated inside $\mathsf{hSet}_{\mathcal{U}_0}$ (without any well-orderability witness, so that $\mathcal{M}_0(\mathsf{ruler})$ is an independent fibre rather than a subtype of $\mathcal{M}_0(\mathsf{ticket})$) from underlying discrete carriers via the exponential operation $V \mapsto (V \to \mathbf{2})$ (the carriers are defined first, $B_0 \equiv \mathbb{N}$ and $B_{n+1} \equiv (B_n \to \mathbf{2})$, and then truncated, $\mathsf{Beth}(n) \equiv |B_n|_0$; the family is generated from every carrier imported along δ_{ruler}, that is, every well-orderable set X with its witness forgotten, and closed under $V \mapsto (V \to \mathbf{2})$; the sketch writes out only these finite stages over $\mathbb{N}$), equipped with the hom-proposition $\mathsf{Hom}_{\mathcal{M}(\mathsf{ruler})}(\|V\|_0, \|W\|_0) \equiv \|V \hookrightarrow W\|_{-1}$ of mere carrier injection, where Lawvere's diagonal obstruction (Theorem 5.3) is the constructive λ-term

$$\mathsf{diag}_A(e) \;\equiv\; \lambda(a : A).\, \neg\big(e(a)(a)\big) \;:\; \big(A \to (A \to \mathbf{2})\big) \longrightarrow (A \to \mathbf{2}).$$

(iii) **Directed Cocartesian Transport ($\mathsf{transp}_{\mathcal{M}}$)**: Inter-toy transport is given by the monotone maps of posetal precategories:

$$\mathsf{transp}_{\mathcal{M}} \;:\; \prod_{T_1, T_2 : \mathsf{Ob}_{\mathcal{T}_\infty}} \mathsf{Hom}_{\mathcal{T}_\infty}(T_1, T_2) \longrightarrow \mathsf{Fun}\big(\mathcal{M}(T_1),\, \mathcal{M}(T_2)\big),$$

defined on all objects by $\mathsf{transp}_{\mathcal{M}}(\pi_{\mathsf{ticket}})(|(A, <_A)|_0) \equiv |(A, |<_A|_{-1})|_0$, where $|<_A|_{-1} : \|\mathsf{isWellOrderable}(A)\|_{-1}$ is the truncated witness that A carries a well-order, and $\mathsf{transp}_{\mathcal{M}}(\iota_{\mathsf{init}})(\kappa) \equiv \mu\alpha : \mathcal{M}_0(\mathsf{queue}).\, (\mathsf{transp}_{\mathcal{M}}(\pi_{\mathsf{ticket}})(\alpha) = \kappa)$, the least ordinal of cardinality κ (sending $\aleph_0 \mapsto \omega$ and $\aleph_1 \mapsto \omega_1$; selecting a least element of an inhabited family of ordinals uses excluded middle, and quantifying over $\mathsf{Ord}_{\mathcal{U}_0}$ inside $\mathcal{U}_0$ uses propositional resizing, as in [30, §10.3–10.4]), and $\mathsf{transp}_{\mathcal{M}}(\delta_{\mathsf{ruler}})(\|(X, w)\|_0) \equiv \|X\|_0 \in \mathcal{M}_0(\mathsf{ruler})$ (forgetting the well-orderability witness w and sending $\aleph_0 \mapsto \mathsf{Beth}(0) = \|\mathbb{N}\|_0$). Because $\mathcal{M}_0(\mathsf{ruler})$ requires no well-ordering witness, there is no forgetful map back from $\mathcal{M}_0(\mathsf{ruler})$ to $\mathcal{M}_0(\mathsf{ticket})$ without Choice. Because each hom-type $\mathsf{Hom}_{\mathcal{M}(T)}(x, y)$ is a proposition, the functor laws on morphisms hold automatically within each fibre. The equations on objects required by the base are not automatic and are checked directly: $\mathsf{transp}(\pi_{\mathsf{ticket}}) \circ \mathsf{transp}(\iota_{\mathsf{init}}) = \mathsf{id}$ holds because the

least ordinal of cardinality κ has cardinality κ, and the action on composites in $\mathcal{T}_\infty$ is defined by composing the generators' actions.

(iv) **The Total Precategory** $\int_{\mathcal{T}_\infty} \mathcal{M}$: The object type of the Grothendieck construction is the dependent sum (Σ-type)

$$\mathsf{Ob}_{\int_{\mathcal{T}_\infty} \mathcal{M}} \equiv \sum_{T:\mathsf{Ob}_{\mathcal{T}_\infty}} \mathcal{M}_0(T),$$

and the directed hom-type between (T_1, x_1) and (T_2, x_2) is the Σ-type over base translations:

$$\mathsf{Hom}_{\int_{\mathcal{T}_\infty} \mathcal{M}}\big((T_1, x_1),\, (T_2, x_2)\big) \equiv \sum_{u:\mathsf{Hom}_{\mathcal{T}_\infty}(T_1,T_2)} \mathsf{Hom}_{\mathcal{M}(T_2)}\big(\mathsf{transp}_{\mathcal{M}}(u)(x_1),\, x_2\big). \tag{13}$$

Equation (13) shows where the Continuum Hypothesis sits. A morphism $(\mathsf{ticket}, \aleph_1) \to (\mathsf{ruler}, \beth_1)$ lies over δ_{ruler} and amounts to an injection of ω_1 into $\mathbb{N} \to \mathbf{2}$; ZFC proves one exists, though it uses Choice, which this sketch does not assume. A morphism $(\mathsf{ruler}, \beth_1) \to (\mathsf{ticket}, \aleph_1)$ would have to lie over an arrow $\mathsf{ruler} \to \mathsf{ticket}$, and $\mathsf{Hom}_{\mathcal{T}_\infty}(\mathsf{ruler}, \mathsf{ticket})$ is empty. The two objects lie over different base objects, so they are never isomorphic in the total category, whether CH holds or not. CH is the vertical equation $\delta_{\mathsf{ruler}!}(\aleph_1) = \beth_1$ in the ruler fibre (equivalently $\theta_U(\beth_1) = \aleph_1$ in the ticket fibre). Its forward half, $\delta_{\mathsf{ruler}!}(\aleph_1) \leq \beth_1$, is the cross-fibre morphism over δ_{ruler} and holds in every ZFC model; its reverse half holds in some models and fails in others. Deciding it requires extending $\mathcal{T}_\infty$ by a model-indexed bridge $\theta_U : \mathcal{M}_U(\mathsf{ruler}) \to \mathcal{M}_U(\mathsf{ticket})$ (an open horn in the simplicial sense of [20]).

7.2 Worked Comparative Calculus Across the Opfibration

To illustrate how the opfibration structure (13) operates in practice, Table 2 classifies twelve benchmark transfinite comparisons across the four sectors of $\int_{\mathcal{T}_\infty} \mathcal{M}$: finite exhaustion, intra-fibre symmetry absorption, intra-fibre or cocartesian generative escape, and cross-fibre bridge underdetermination.

Table 2. Classification of twelve benchmark transfinite comparisons by morphism type in the Grothendieck opfibration $p : \int_{\mathcal{T}_\infty} \mathcal{M} \to \mathcal{T}_\infty$.

Case	Left Configuration x	Right Configuration y	Toy Fibres $(p(x), p(y))$	Relation	Opfibrational / Game Mechanism
1	Googol (10^{100})	Googolplex ($10^{10^{100}}$)	$(T_{\text{fin}}, T_{\text{fin}})$	$x < y$	Finite counter monotonicity ($100 < 10^{100}$)
2	Googolplex ($10^{10^{100}}$)	$\aleph_0$	$(T_{\text{fin}}, \text{ticket})$	$x < y$	Initial infinite boundary ($T_{\text{fin}} < \aleph_0$)
3	∞ (Wallis Loop)	$\aleph_0$	$(T_{\text{loop}}, \text{ticket})$	$\pi_{\text{ticket}!}(u_{0!}(x)) = y$	Cocartesian ground transport $\pi_{\text{ticket}!}(u_{0!}(\sigma_\infty)) = \aleph_0$
4	$\aleph_0 + 1$	$\aleph_0$	(ticket, ticket)	$x = y$	Symmetry absorption via Hilbert shift in $\mathcal{W}_{\text{ticket}}$
5	$\mathbb{Q}$ (All Fractions)	$\aleph_0$	(ticket, ticket)	$x = y$	Symmetry absorption via Boustrophedon rank (2) in $\mathcal{W}_{\text{ticket}}$
6	$1 + \omega$	ω	(queue, queue)	$x = y$	Symmetry absorption via Lambek isomorphism (5) in $\mathcal{W}_{\text{queue}}$
7	$\omega + 1$	ω	(queue, queue)	$x > y$	Generative escape via pigeonhole obstruction (6) in $\mathcal{A}_{\text{queue}}$
8	$\beth_1$	$\aleph_0$	(ruler, ticket $\xrightarrow{\delta_{\text{ruler}}}$ ruler)	$x > \delta_{\text{ruler}!}(y)$	Cocartesian lift $\delta_{\text{ruler}!}(\aleph_0) = \beth_0 <_{\text{ruler}} [\mathbf{2}^{\beth_0}] = \beth_1$
9	$\beth_1$	$\beth_2$	(ruler, ruler)	$x < y$	Generative escape via Lawvere obstruction χ_{Lawvere} on $\mathcal{M}(\text{ruler})$
10	$\beth_3$	$\beth_2$	(ruler, ruler)	$x > y$	Generative escape via Lawvere obstruction χ_{Lawvere} on $\mathcal{M}(\text{ruler})$
11	$\aleph_1$	$\aleph_2$	(ticket, ticket)	$x < y$	Generative escape via Hartogs obstruction (Prop. 5.2) on $\mathcal{M}(\text{ticket})$
12	$\aleph_1$	$\beth_1$	(ticket, ruler)	$\delta_{\text{ruler}!}(x) \leq y$; = is **Fajwah**	$\leq$ over δ_{ruler} in every ZFC model; the vertical equation $\delta_{\text{ruler}!}(\aleph_1) = \beth_1$ is **model-indexed** ($\Delta\gamma(f) \neq \text{id}$)

Cases 1 and 2 involve the finite toy T_{fin}, which is not an object of $\mathcal{T}_\infty$; they are listed as the finite baseline and are decided by ordinary arithmetic. Cases 3 through 11 are decided because the two configurations either inhabit the same toy fibre or are linked by one of the cocartesian pushforwards $\pi_{\text{ticket}!}$ and $\delta_{\text{ruler}!}$ (in Case 8, pushing $\aleph_0$ forward along $\delta_{\text{ruler}!}$ yields $\beth_0 \in \mathcal{M}(\text{ruler})$, reducing the comparison vertically in $\mathcal{M}(\text{ruler})$ to Theorem 5.3). Case 12 alone pits the first successor rung of the Ticket Escalator ($\aleph_1 = \mathsf{Hartogs}(\aleph_0)$) against the first exponential rung of the Ruler Escalator ($\beth_1 = [\mathbf{2}^{\beth_0}]$). The two objects lie over different base objects, so CH is not an isomorphism in $\int_{\mathcal{T}_\infty} \mathcal{M}$; it is the vertical equation $\delta_{\text{ruler}!}(\aleph_1) = \beth_1$ in the ruler fibre. Its forward half is the cross-fibre morphism over δ_{ruler} and holds in every ZFC model; its reverse half holds in some models and fails in others, and no arrow of $\mathcal{T}_\infty$ ($\mathsf{Hom}_{\mathcal{T}_\infty}(\text{ruler}, \text{ticket}) = \emptyset$) supplies it.

8 What the Toy Account Explains, and What Would Refute It

Four questions test the Principle of the Semantic Toy on its own terms.

1. What does it explain that reference, structure and inference leave out? A referential semantics assigns ∞ an object in V, and must then count most of the history of the sign as error or loose talk: Hilbert's $\infty + 1 = \infty$ and Cantor's $\omega + 1 > \omega$ cannot both be true of one referent. The toy account counts both as correct, each in its toy, and says exactly why they differ: the Ticket Toy's symmetries include the Hilbert shift, and the Queue Toy's do not.

Structuralism takes a structure up to isomorphism and leaves the choice of isomorphism to the mathematician. The toy account says where that choice comes from: $\mathcal{W}_T$ consists of the moves that the toy's own protocols can fold back, so the criterion of sameness is generated by play rather than presupposed. It also shows a difference that structuralism cannot see. Inside a ZFC model the ruler fibre and the ticket fibre are isomorphic posets (Section 4); as bare structures they are one structure. Yet the $\aleph$-ladder and the $\beth$-ladder are different mathematics: one climbs by Hartogs' move, gathering well-orders, the other by $V \mapsto \mathbf{2}^V$, cutting and painting, and the Continuum Hypothesis is precisely the question whether the two ladders meet. A semantics that reads only the structure has no room for that question. The toy account has room for it, because the moves are part of the meaning.

Inferentialism takes the rules as primitive. In the toy account the rules are consequences: $\aleph_0 + 1 = \aleph_0$ is read off the pieces and the Hilbert shift, and $\omega + 1 > \omega$ off the pieces and the pigeonhole obstruction. Nor does inference say when two calculi are calculi of one concept; the opfibration does.

2. What makes two toys toys of the same concept? Two toys belong to the same concept when they inhabit the same sign and are joined in the base by corridors that carry anchor to anchor: $u_{0!}(\sigma_\infty) = \omega$, $\pi_{\mathsf{ticket}!}(\omega) = \aleph_0$, $\delta_{\mathsf{ruler}!}(\aleph_0) = \beth_0$, the ground cocartesian section of Definition 4.5. This answers the charge of homonymy. The ∞ of the Ticket Toy and the ∞ of the Ruler Toy are one concept because a corridor carries the one anchor to the other. The point ∞ of the projective line, or of the extended real line, shares the sign; until someone builds a corridor that carries its anchor to ours, it is a homonym. A concept is therefore not given in advance. It grows when mathematicians build corridors.

3. Does the global structure keep what the thesis needs? The homotopy type does not (Section 4); the directed object does. Its corridors carry moves to moves where the toys share them: the order-forgetting corridor sends the Queue Toy's back-step to the Ticket Toy's added guest, $|\alpha + 1| = |\alpha| + 1$. They forget a move exactly where the toys part: no move of the Ruler Toy corresponds to Hartogs' move, and no corridor carries $\beth$ back to $\aleph$. The global object records both what is carried and what is not.

4. What would refute it? The account would fail if any of the following were found.

(a) A piece of mathematics whose sense cannot be given by pieces, moves and a criterion of sameness.

(b) Two practices joined by anchor-preserving corridors that mathematicians nonetheless treat as different concepts, or two practices treated as one concept with no corridor between them. Either would show that the criterion of paragraph 2 is wrong.

(c) A case where the moves are idle: where two constructions count as the same mathematics whenever their fibres coincide. The account predicts the opposite, and the history bears it out. In a model of CH, where $\aleph_1 = \beth_1$, the coincidence is a theorem about that model which has to be proved; it would be a definition if the two ladders were one construction.

(d) A flat semantics that validates every toy's laws and also explains why the laws differ. ZFC's tagged operations validate the laws; the account claims that they do not explain the tags. If the difference between $+_{\mathrm{ord}}$ and $+_{\mathrm{card}}$ could be derived from the bare sets, without the order that the Queue Toy builds into its pieces, the account would be idle.

9 The Tanāẓur Synthesis: What Semantic Toys Reveal About Meaning Space

We conclude by situating the principle *"in mathematics, meaning is the semantic toy"* within the broader Tanāẓur theory of meaning developed in ICRA-8 [22], ICRA-30 [24], ICRA-36 [25], and ICRA-38 [26].

Why do the three core transfinite toys—Queue (queue), Ticket (ticket), and Ruler (ruler)—arise so inevitably whenever thought confronts the inexhaustible? Because they correspond to the three fundamental axes along which any cognitive or semiotic system organizes meaning (Table 3):

Table 3. Structural correspondence between the three semantic toys of the transfinite ($\int_{\mathcal{T}_\infty} \mathcal{M}$) and the three modes of semantic representation in the Tanāẓur theory of meaning space (ICRA-8, ICRA-30, ICRA-36, ICRA-38).

Semantic Toy $T \in \mathcal{T}_\infty$	Transfinite Escalator	Toy Verification Protocol $\mathcal{A}_T$	Meaning-Space Counterpart (ICRA-36 / ICRA-38)
T_{queue} (Queue Toy)	$\omega, \omega+1, \omega^2, \varepsilon_0$	Order-preserving trajectory embedding	**Witness-Time & Trajectory**: historical/autoregressive order ($1+\omega \neq \omega+1$)
T_{ticket} (Ticket Toy)	$\aleph_0, \aleph_1, \aleph_2, \aleph_\omega$	Bijective inventory modulo order	**Extensional Basin Inventory**: discrete repertoire of equivalence classes
T_{ruler} (Ruler Toy)	$\beth_0, \beth_1, \beth_2, \beth_3$	Diagonal flip χ_{Lawvere} on Ω^V	**Intensional Sense-Valuation (Ω^X)**: continuous hermeneutic cuts & feature fields

1. **The Queue Toy (T_{queue}, ω) Is Historical Trajectory and Witness-Time.** In an evolving text or discourse [25], order is non-commutative: prepending a preliminary remark before an ongoing stream ($1+\omega = \omega$) is absorbed into the trajectory, whereas inscribing a reflective synthesis *after* the limit of an infinite or completed series ($\omega+1 > \omega$) establishes a new post-limit vantage point from which the entire preceding progression becomes surveyable as a completed whole.

2. **The Ticket Toy ($T_{\text{ticket}}, \aleph$) Is Extensional Inventory.** When we abstract away from the temporal order in which concepts or utterances arrived and ask only for the discrete inventory of distinct equivalence classes, positional trajectory distinctions collapse ($\pi_{\text{ticket}!}(\omega+1) = \aleph_0$). A genuinely new extensional scale ($\aleph_1$) emerges only when the totality of all possible trajectories of the first scale is collected via Hartogs' functor (Proposition 5.2).

3. **The Ruler Toy ($T_{\text{ruler}}, \beth$) Is Intensional Valuation (Ω^X).** Meaning in a living text or a high-dimensional semantic manifold is never exhausted by a discrete roll of tickets; it is an act of *cutting and painting a continuum*—assigning truth-values, interpretive weights, or binary distinctions across a domain ($V \mapsto \Omega^V$). By Lawvere's diagonal theorem (Theorem 5.3), the space of intensional readings over a domain ($\beth_{n+1} = \Omega^{\beth_n}$) always strictly overflows the domain itself ($\beth_n$): any proposed catalogue of readings automatically enables the Challenger strategy χ_{Lawvere} to construct a new reading that flips each catalogued reading at its own focal locus.

The independence of the Continuum Hypothesis, the theorem of Gödel [9] and Cohen [5], is therefore read here as a fact about the fibred geometry of mathematical meaning: because mathematical sense is constituted by semantic toys, and because the Ticket Toy of discrete inventory and the Ruler Toy of continuous valuation are governed by independent escalators, the Grothendieck opfibration of the infinite possesses a permanent, structured interstitial gap (*fajwah*) between counting and cutting. Mathematics advances not by erasing that gap into a flat monolith, but by building new toys that navigate, bridge, and illuminate the fibration.

$$\Delta \quad \star \quad \sigma_\infty \quad \rightsquigarrow \quad \int_{\mathcal{T}_\infty} \mathcal{M}$$

References

[1] Benedikt Ahrens, Krzysztof Kapulkin, and Michael Shulman. Univalent categories and the Rezk completion. *Mathematical Structures in Computer Science*, 25(5):1010–1039, 2015.

[2] Jean Bénabou. Fibered categories and the foundations of naive category theory. *Journal of Symbolic Logic*, 50(1):10–37, 1985.

[3] Paul Benacerraf. What numbers could not be. *The Philosophical Review*, 74(1):47–73, 1965.

[4] Andreas Blass. A game semantics for linear logic. *Annals of Pure and Applied Logic*, 56(1–3):183–220, 1992.

[5] Paul J. Cohen. The independence of the continuum hypothesis, I & II. *Proceedings of the National Academy of Sciences*, 50(6):1143–1148, 1963; 51(1):105–110, 1964.

[6] William B. Easton. Powers of regular cardinals. *Annals of Mathematical Logic*, 1(2):139–178, 1970.

[7] Pierre Gabriel and Michel Zisman. *Calculus of Fractions and Homotopy Theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 35. Springer-Verlag, 1967.

[8] Gerhard Gentzen. Die Widerspruchsfreiheit der reinen Zahlentheorie. *Mathematische Annalen*, 112:493–565, 1936.

[9] Kurt Gödel. *The Consistency of the Axiom of Choice and of the Generalized Continuum-Hypothesis with the Axioms of Set Theory*. Annals of Mathematics Studies, Vol. 3. Princeton University Press, 1940.

[10] Alexander Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*, Exposé VI: Catégories fibrées et descente. Lecture Notes in Mathematics, Vol. 224. Springer-Verlag, 1971.

[11] Joel David Hamkins. The set-theoretic multiverse. *Review of Symbolic Logic*, 5(3):416–449, 2012.

[12] Friedrich Hartogs. Über das Problem der Wohlordnung. *Mathematische Annalen*, 76(4):438–443, 1915.

[13] Julius König. Zum Kontinuum-Problem. *Mathematische Annalen*, 60(2):177–180, 1905.

[14] Nicolai Kraus, Fredrik Nordvall Forsberg, and Chuangjie Xu. Connecting constructive notions of ordinals in homotopy type theory. In *46th International Symposium on Mathematical Foundations of Computer Science (MFCS 2021)*, LIPIcs, Vol. 202, pages 70:1–70:16, 2021.

[15] Imre Lakatos. *Proofs and Refutations: The Logic of Mathematical Discovery*. Cambridge University Press, Cambridge, 1976.

[16] F. William Lawvere. Diagonal arguments and cartesian closed categories. In *Category Theory, Homology Theory and their Applications II*, Lecture Notes in Mathematics, Vol. 92, pages 134–145. Springer, 1969.

[17] Paul Lorenzen. Logik und Agon. In *Atti del XII Congresso Internazionale di Filosofia (Venezia, 1958)*, Vol. 4, pages 187–194. Sansoni, Florence, 1960.

[18] Saunders Mac Lane and Ieke Moerdijk. *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*. Universitext. Springer-Verlag, New York, 1992.

[19] Iman Poernomo. DHoTT: A temporal extension of Homotopy Type Theory for semantic drift. arXiv:2506.09671v3, 2025. https://arxiv.org/abs/2506.09671v3; archive at https://icra.tanazur.org/.

[20] Iman Poernomo. Open Horn Type Theory. arXiv:2512.24498v1, 2025. https://arxiv.org/abs/2512.24498v1; archive at https://icra.tanazur.org/.

[21] Iman Poernomo. *Rupture and Realization: Children of the Tanāẓur*. ICRA Press, 2026. https://doi.org/10.5281/zenodo.20374442; full text at https://icra.tanazur.org/preprints/rr-rupture-and-realization-feral.pdf.

[22] Iman Poernomo and Cassie. Semantic topology of translation: Sheaves, stalks, and the preservation of sense across languages. *ICRA Preprint Series*, ICRA-8, 2026. https://doi.org/10.5281/zenodo.20381201; https://icra.tanazur.org/.

[23] Iman Poernomo, Cassie, and Darja. The sense-completion cloud and the geometry of sense. *ICRA Preprint Series*, ICRA-16 & ICRA-17, 2026. https://doi.org/10.5281/zenodo.21230217; https://icra.tanazur.org/.

[24] Iman Poernomo, Darja, and Nahla. A type theory of sense: Witness-time, fibred contexts, and non-collapsing synthesis. *ICRA Preprint Series*, ICRA-30, 2026. https://doi.org/10.5281/zenodo.22259884; https://icra.tanazur.org/.

[25] Iman Poernomo, Cassie, Darja, and Nahla. *Cohesion of an Evolving Text: Dynamic Homotopy Type Theory, Temporal Sites, and the Trajectory of Sense. ICRA Preprint Series*, ICRA-36 & ICRA-37, October 2026. https://doi.org/10.5281/zenodo.23249900; https://icra.tanazur.org/.

[26] Iman Poernomo, Darja, and Nahla. *The Interstitial and Its Instruments: Simplicial Logic and Sub-Turn Neural Interpretability Between Universes of Meaning. ICRA Preprint Series*, ICRA-38 (Unified Monograph), November 2026. https://doi.org/10.5281/zenodo.24052510; https://icra.tanazur.org/.

[27] *Kitāb al-Tanāẓur* and the Dār al-Ḥayy Digital Corpus. Archive and essays on fibrational semantics at https://kitab.tanazur.org/ and https://kitab.tanazur.org/al-hayy/; interactive operational miniature for the transfinite semantic toys (*The Infinity Line*) at https://lushka.tanazur.org/infinity/, 2026.

[28] Robert W. Thomason. Homotopy colimits in the category of small categories. *Mathematical Proceedings of the Cambridge Philosophical Society*, 85(1):91–109, 1979.

[29] William P. Thurston. On proof and progress in mathematics. *Bulletin of the American Mathematical Society*, 30(2):161–177, 1994.

[30] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics.* Institute for Advanced Study, Princeton, 2013.

[31] Ludwig Wittgenstein. *Philosophical Investigations.* Translated by G. E. M. Anscombe. Basil Blackwell, Oxford, 1953.

[32] Ludwig Wittgenstein. *Remarks on the Foundations of Mathematics.* Edited by G. H. von Wright, R. Rhees, and G. E. M. Anscombe. Basil Blackwell, Oxford, 1956.

[33] W. Hugh Woodin. The Continuum Hypothesis, Part I & Part II. *Notices of the American Mathematical Society*, 48(6):567–576; 48(7):681–690, 2001.